



Modeling and Analysis of a Fractional Visceral Leishmaniasis with Caputo and Caputo–Fabrizio derivatives

Dalal Khalid Almutairi^a, Mohamed A. Abdoon^b, Salih Yousuf Mohamed Salih^b, Shahinaz A. Elsamani^b, Fathelrhman EL Guma^c, Mohammed Berir^{b,c,*}

^aDepartment of Mathematics, College of Education (Majmaah), Majmaah University, P.O.Box 66, Al-Majmaah, 11952, Saudi Arabia.

^bDepartment of Mathematics, Faculty of Science, Bakht Al-Ruda University, Duwaym, Sudan.

^cDepartment of Mathematics, Faculty of Science and Arts in Baljurashi, Albaha University, Albaha, Saudi Arabia.

Abstract

Visceral leishmaniasis is one recent example of a global illness that demands our best efforts at understanding. Thus, mathematical modeling may be utilized to learn more about and make better epidemic forecasts. By taking into account the Caputo and Caputo-Fabrizio derivatives, a frictional model of visceral leishmaniasis was mathematically examined based on real data from Gedaref State, Sudan. The stability analysis for Caputo and Caputo-Fabrizio derivatives is analyzed. The suggested ordinary and fractional differential mathematical models are then simulated numerically. Using the Adams-Bashforth method, numerical simulations are conducted. The results demonstrate that the Caputo-Fabrizio derivative yields more precise solutions for fractional differential equations.

DOI:10.46481/jnsps.2023.1453

Keywords: Leishmaniasis, Modelling, Caputo, Caputo–Fabrizio, Sudan.

Article History :

Received: 16 March 2023

Received in revised form: 21 May 2023

Accepted for publication: 19 June 2023

Published: 26 July 2023

© 2023 The Author(s). Published by the Nigerian Society of Physical Sciences under the terms of the Creative Commons Attribution 4.0 International license (<https://creativecommons.org/licenses/by/4.0>). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Communicated by: B. J. Falaye

1. Introduction

Visceral leishmaniasis, or kala azar, is a lethal vector-borne illness. India, Bangladesh, and Nepal have achieved substantial headway in lowering VL cases. East Africa has made less progress, especially with South Sudan's continuous endemicity and VL outbreaks during the past 40 years. Lack of infrastructure, clinical staff, IDPs, and hunger have hampered VL management in, and longer-term hazards to diagnostic kits and medications. Pentavalent antimonials have been the backbone

of VL treatment for decades, and resistance to them, as previously demonstrated in the Indian subcontinent, provides another major barrier to VL treatment and management. To prevent monotherapy and minimize treatment duration, first-line 30-day sodium stibogluconate SS is substituted with a 17-day injectable combination regimen of SSG and PM in WHO recommendations in 2010 and Sudan Ministry of Health guidelines in 2011. Since 2012, AmBisome has been donated to WHO for these purposes. In East Africa, SSG/PM combo treatment had a 5% recurrence rate. Relapse may be due to insufficient cellular immunity following therapy due to HIV, TB, or malnutrition, or inadequate treatment resulting in considerable chronic parasitaemia after initial clinical cure. In places like Sudan, where

*Corresponding author

Email address: midriss@bu.edu.sa (Mohammed Berir)

active patient follow-up is difficult and not common, VL recurrence rates are passively evaluated by VL re-treatment admissions as a proportion of overall VL admissions. Passive monitoring shows an increase in re-treatment rates in recent years [1-7].

In the last 30 years, fractional calculus and nonlinear equations has become more well-known and important. Fractional differential equations are used in physics, chemical engineering, mathematical biology, and finance [8-16]. Simulating a fractional model simultaneously with a Caputo derivative and a CF derivative. In addition, modeling and graphing with the fractional derivative is a highly effective technique for demonstrating leishmaniasis using MATLAB. This could be done to better comprehend the infection. Using fractional derivatives as a research strategy for natural occurrences may result in more precise findings than other methods. As a result of this model's use of a non-singleton kernel, the CF derivative has significantly improved predictive abilities.

2. Preliminaries

Definition 2.1. Riemann-Liouville fractional integral (RLI) operator of order $\alpha > 0$ for a function $y(\tau)$ is given by [17]:

$$D^\alpha y(t) = \frac{1}{\Gamma(n-\alpha)} \int_0^t (t-\tau)^{n-\alpha-1} y^n(\tau) d\tau = I^{n-\alpha} y^n(t), t > 0. \tag{1}$$

Definition 2.2. For $y \in H^1(0, t), t > 0, T > 0, \alpha \in (0, 1]$ Then the CF fractional operator [17] is given by

$${}^C_0 D_t^\alpha y(t) = \frac{B(\alpha)}{1-\alpha} \frac{d}{dt} \int_0^t y(\tau) e^{-\alpha \frac{t-\tau}{1-\alpha}} d\tau, 0 < \alpha < 1. \tag{2}$$

In this expression $B(\alpha)$ satisfies the condition $B(0) = B(1) = 1$.

Definition 2.3. Caputo derivative of order $0 \leq n-1 < \alpha < n$ with the lower limit zero for a function $y(\tau)$ is given by [18]:

$$I^\alpha y(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} y(\tau) d\tau, t > 0. \tag{3}$$

3. Anthropologic Visceral Leishmaniosis model with Caputo derivative

In this Section, we describe the leishmaniasis model, which includes four subpopulations: susceptible, infectious, Recovered, and Recovered with permanent immunity, for the human population, and two compartments for the reservoir population: susceptible and infected. In addition to that, we have two compartments for sandflies: susceptible and infected. The human population is the only population in the model that has permanent immunity. The positivity, reproduction number, and equilibrium solutions of the model that was established in this work

have all been determined to be free of leishmaniasis. Additionally, the leishmaniasis cases, along with their respective localities and global stability properties, have also been determined. We obtain the model formulation by using a new variable:

$$s_h(t) = \frac{S_H}{N_H}, i_h(t) = \frac{I_H}{N_H}, p_h(t) = \frac{P_H}{N_H}, r_h(t) = \frac{R_H}{N_H}, s_r(t) = \frac{S_R}{N_R}, I_r(t) = \frac{I_R}{N_R}, s_v(t) = \frac{S_V}{N_V}, i_v(t) = \frac{I_V}{N_V}, s_h(t) = \frac{S_H}{N_H}, m = \frac{N_V}{N_H} \text{ and } N = \frac{N_V}{N_R}.$$

The system of differential equations is given by:

$$\begin{cases} {}^C_0 D_t^\alpha i_h = abmi_v N_h - (\alpha_1 + \delta + \frac{A_H}{N_H} - \delta i_h) i_h, \\ {}^C_0 D_t^\alpha p_h = (1 - \sigma) \alpha_1 i_h - (\alpha_2 + \beta + \frac{A_H}{N_H} - \delta i_h) p_h, \\ {}^C_0 D_t^\alpha i_r = abni_v s_r - \frac{A_H}{N_H} i_r, \\ {}^C_0 D_t^\alpha i_v = aci_h S_v + acp_h S_v + aci_r S_v - \frac{A_V}{N_V} i_v, \\ {}^C_0 D_t^\alpha s_h = \frac{A_H}{N_H} - [abmi_v + \frac{A_H}{N_H} - \delta i_h] s_h, \\ {}^C_0 D_t^\alpha r_h = \sigma \alpha_1 i_h + (\alpha_2 + \beta) P_h - [\frac{A_H}{N_H} - \delta i_h] r_h, \\ {}^C_0 D_t^\alpha S_r = \frac{A_R}{N_R} - abni_v s_r - \frac{A_H}{N_H} S_r, \\ {}^C_0 D_t^\alpha s_v = \frac{A_V}{N_V} - [aci_h + acP_h + \frac{A_V}{N_V}] s_v, \end{cases} \tag{4}$$

with initial conditions:

$$s_h(0) = c_1, i_h(0) = c_2, r_h(0) = c_3, s_r(0) = c_4, I_r(0) = c_5, s_v(0) = c_6, i_v(0) = c_7.$$

4. Anthropologic Visceral Leishmaniosis model with FC derivative

In this Section, we obtain the fractional model formulation under Caputo-Fabrizio derivatives: $s_h(t) = \frac{S_H}{N_H}, i_h(t) = \frac{I_H}{N_H}, p_h(t) = \frac{P_H}{N_H}, r_h(t) = \frac{R_H}{N_H}, s_r(t) = \frac{S_R}{N_R}, I_r(t) = \frac{I_R}{N_R}, s_v(t) = \frac{S_V}{N_V}, i_v(t) = \frac{I_V}{N_V}, s_h(t) = \frac{S_H}{N_H}, m = \frac{N_V}{N_H}$ and $N = \frac{N_V}{N_R}$ The system of differential equations is given by:

$$\begin{cases} {}^{FC}_0 D_t^\alpha i_h = abmi_v N_h - (\alpha_1 + \delta + \frac{A_H}{N_H} - \delta i_h) i_h, \\ {}^{FC}_0 D_t^\alpha p_h = (1 - \sigma) \alpha_1 i_h - (\alpha_2 + \beta + \frac{A_H}{N_H} - \delta i_h) p_h, \\ {}^{FC}_0 D_t^\alpha i_r = abni_v s_r - \frac{A_H}{N_H} i_r, \\ {}^{FC}_0 D_t^\alpha i_v = aci_h S_v + acp_h S_v + aci_r S_v - \frac{A_V}{N_V} i_v, \\ {}^{FC}_0 D_t^\alpha s_h = \frac{A_H}{N_H} - [abmi_v + \frac{A_H}{N_H} - \delta i_h] s_h, \\ {}^{FC}_0 D_t^\alpha r_h = \sigma \alpha_1 i_h + (\alpha_2 + \beta) P_h - [\frac{A_H}{N_H} - \delta i_h] r_h, \\ {}^{FC}_0 D_t^\alpha S_r = \frac{A_R}{N_R} - abni_v s_r - \frac{A_H}{N_H} S_r, \\ {}^{FC}_0 D_t^\alpha s_v = \frac{A_V}{N_V} - [aci_h + acP_h + \frac{A_V}{N_V}] s_v, \end{cases} \tag{5}$$

with initial conditions: $s_h(0) = c_1, i_h(0) = c_2, r_h(0) = c_3, s_r(0) = c_4, I_r(0) = c_5, s_v(0) = c_6, i_v(0) = c_7.$

5. Stability analysis

In this part we discuss the stability of epidemiological model, the equilibrium points, eigenvalues value and the Jacobian matrix for the model (1).

Table 1: Description of the variables for model.

Variable	Description
$N_H(t)$	Human host population
$N_R(t)$	Reservoir host population
$N_V(t)$	Vector population
$S_H(t)$	Susceptible humans
$P_H(t)$	Recovered and have permanent immunity
$I_H(t)$	Infected humans
$R_H(t)$	Recovery humans
$R_s(t)$	Susceptible reservoir
$I_R(t)$	Infected reservoir
$S_V(t)$	Susceptible sandflies
$I_V(t)$	Infected sandflies

Table 2: Parameters values of the leishmaniasis model.

Parameter	Description	Value	Source
a	Biting rate of sandflies	0.2856 day ⁻¹	[16]
b	Progression rate of VL in sandfly	0.22 day ⁻¹	[16]
c	Progression rate of VL in human and reservoir	0.0714 day ⁻¹	[16]
A_H	Human recruitment rate	10.1009 day ⁻¹	Estimated
A_R	Reservoir recruitment rate	19.7795 day ⁻¹	Estimated
A_V	Vector recruitment rate	38858.62 day ⁻¹	Estimated
μ_h	Natural mortality rate of humans	4.341e - 6 day ⁻¹	[2]
μ_r	Natural mortality rate of reservoirs	0.0017 day ⁻¹	[1]
μ_v	Natural mortality rate of vectors	0.0668 day ⁻¹	[1]
α_1	Treatment rate of VL	0.02	[2]
α_2	PKDL recovery rate without treatment	0.033	[20]
σ	Recovery rate from VL infection after treatment	0.9	[1]
1 - σ	Developing PKDL rate after treatment	0.1	[1]
δ	Death rate due to VL	0.011	[16]
β	PKDL recovery rate after treatment	0.9	[1]

5.1. Equilibria

The equilibrium points of dynamics (5) are computed solving the nonlinear system.

Table 3: The equilibrium points of the system.

E_i	Equilibria
E_1	(0, 0, 0, 0, 1, 0, 711.58, 1)
E_2	(32.5436, 0.1132, 711.5801, 22.7897, -29.7254, -1.9314, 0, 0)
E_3	(-31.1459, -0.0488, 711.58, -21.8109, 33.9641, -1.7693, 0, 0)
E_4	(85.0303, -72.8759, 711.5801, 59.5451, -82.2121, -71.0578, 0, 0)
E_5	(2.8182, 0.0062, -2.8244, 0, 0, -1.8244, 714.4046, 0)
E_6	(2.8182, 0.0062, 711.5801, 252.9373, 0, -1.8244, 0, 0)
E_7	(0, 0, 0, 0, -7.2892e11, 7.2892e11, 0, 0)
E_8	(0, 0, 0, 5.1155e8, 0, 0, 0, 0)

5.2. The Jacobian matrix for the model:

Here, we talk about this epidemiological model stability. The disease-free equilibrium point is given as $E_1 = (0, 0, 0, 0, 1, 0, 711.58, 1)$ and the endemic equilibrium points $E_8 = (0, 0, 0, 5.1155e8, 0, 0, 0, 0)$.

$$J(E_1) = \begin{bmatrix} -0.031 & 0 & 0 & 0.0157 & 0 & 0 & 0 & 0 \\ 0.002 & -0.933 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -4.297e^{-8} & 15907.35 & 0 & 0 & 0 & 0 \\ 0.02 & 0.02 & 0.02 & -3.438e^{-11} & 0 & 0 & 0 & 0 \\ 0.011 & 0 & 0 & -0.0157 & -4.297e^{-8} & 0 & 0 & 0 \\ 0.018 & 0.0933 & 0 & 0 & 0 & -4.297e^{-8} & 0 & 0 \\ 0 & 0 & 0 & -15907.35 & 0 & 0 & -4.297e^{-8} & 0 \\ -0.02 & -0.02 & 0 & 0 & 0 & 0 & 0 & -3.438e^{-11} \end{bmatrix} \tag{6}$$

$$J(E_2) = \begin{bmatrix} -0.031 & 0 & 0 & 0 & 8035456.6 & 0 & 0 & 0 \\ 0.002 & -0.933 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -4.297e^{-8} & -0.0006 & 0 & 0 & 11435742383.06 & 0 \\ 0.000002 & 0.00002 & 0.00002 & -3.438e^{-11} & 0 & 0 & 0 & 14.225 \\ 0 & 0 & 0 & 0 & -8035456.6 & 0 & 0 & 0 \\ 0.028 & 0.0933 & 0 & 0 & 0 & -0.00000003 & 0 & 0 \\ 0 & 0 & 0 & 0.0006 & 0 & 0 & -11435742383.06 & 0 \\ -0.00002 & -0.0002 & 0 & 0 & 0 & 0 & 0 & -0.000000027 \end{bmatrix} \tag{7}$$

Table 4: Variable values.

Eigenvalues	Stability
λ (17.833, 0, 0, 0, 0, -17.833)	Unstable
λ^* (-3.438e ⁻¹¹ , -2.8e ⁻⁸ , -2.8e ⁻⁸ , -4.297e ⁻⁸ , -0.31, -0.933, -035456.6, -1.14e ¹⁰)	Stable

5.3. The Basic Reproduction Number

The basic reproduction number is a baseline statistic in epidemiology and is represented by R_0 , which stands for the predicted value of the secondary infections rate per time unit. Using the equation's fractional model (1), we have fours infected classes, rewrite the system of Equation 1 for the susceptible and

infected classes in the general form:

$$\frac{dx}{dt} = f(x) - v(x), \tag{8}$$

where

$$f(x) = \begin{pmatrix} abmi_v s_h \\ 0 \\ abmi_v s_r \\ ac(i_h + p_h + i_r) s_v \end{pmatrix},$$

$$\text{and } v(x) = \begin{pmatrix} (\alpha_1 + \delta + \mu_h) i_h \\ (\alpha_2 + \beta + \mu_h) p_h - (1 - \sigma) \alpha_1 i_h \\ \mu_r i_r \\ \mu_v i_v \end{pmatrix}. \tag{9}$$

librium point is:

$$F = \begin{pmatrix} 0 & 0 & 0 & abm \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & abm \\ ac & ac & ac & 0 \end{pmatrix}, \tag{10}$$

$$\text{and } V = \begin{pmatrix} \alpha_1 + \delta + \mu_h & 0 & 0 & 0 \\ -(1 - \sigma) \alpha_1 & \alpha_2 + \beta + \mu_h & 0 & 0 \\ 0 & 0 & \mu_r & 0 \\ 0 & 0 & 0 & \mu_v \end{pmatrix}$$

we have

$$R_0 = \rho(FV^{-1}) = \sqrt{\frac{ac[\mu_r abm(\alpha_2 + \delta + \mu_h + (1 - \sigma) \alpha_1) + abn(\alpha_1 + \delta + \mu_h)(\alpha_2 + \delta + \mu_h)]}{\mu_v \mu_r (\alpha_1 + \delta + \mu_h)(\alpha_2 + \delta + \mu_h)}}. \tag{11}$$

Lemma 5.1. *The disease-free equilibrium E_0 is locally asymptotically stable if $R_0 < 1$ and unstable if $R_0 > 1$.*

6. Numerical Simulation and Graphical Representations

This section is devoted to finding the approximate solutions of the proposed models (4) and (5) under fractional operators of Caputo and Caputo-Fabrizio, respectively. We simulate our model using some highly reliable numerical techniques. The finite difference scheme for the initial value problem yields the following numerical techniques for the underlying operators:

$${}^c x_{r+1} = x_0 + \frac{(\Delta t)^\omega}{\Gamma(\omega + 1)} \sum_{k=0}^r [(r - k + 1)^\omega - (r - k)^\omega] F(x_k) + O(\Delta t^2),$$

$${}^{CF} x_{r+1} = x_0 + (1 - \delta)F(x_r) + \delta \Delta t \sum_{k=0}^r F(x_k) + O(\Delta t^2), \tag{12}$$

Table 1 shows a description of the variables in the model. Figures 1 and 2 were obtained with the Caputo (4) and CF methods (5) using the parameters in Table 2. Tables 3 & 4 show a summary of equilibrium points and the corresponding eigenvalues of the Jacobian matrix.

7. Conclusion

A fractional model was simulated by using a Caputo derivative as well as a CF derivative simultaneously. In addition, modeling and graphing with the aid of the fractional derivative is a very effective approach that can be used to show leishmaniasis with the use of MATLAB. This may be done in order to better understand the infection. When doing research on natural events, using fractional derivatives as a strategy might lead to more precise findings than other approaches. Due to the fact that this model employs a non-singleton kernel, the CF derivative has much enhanced prediction capabilities. This research was carried out in the hope that it will be a useful resource for future applications and explorations of simulation by using a Caputo derivative, and to investigate new methods such as those in Refs. [18-29]




Figure 1: Systems of fractional orders model for $\alpha=0.99$.




Figure 2: Systems of fractional orders model for $\alpha=1$ (First part)

Now, the Jacobian of $f(x)$ and $v(x)$ of the disease free equi-

References

- [1] E. E. Zijlstra & A. M. el-Hassan, "Leishmaniasis in Sudan. 3. Visceral leishmaniasis", *Transactions of the Royal Society of Tropical Medicine and Hygiene* **95** (2001) 27.
- [2] M. Siddig, H. Ghalib, D.C. Shillington, E.A. Petersen & S. Khidir, "Visceral leishmaniasis in Sudan. Clinical features", *Tropical and Geographical Medicine* **42** (1990) 107.
- [3] M. Ahmed, A. A. Abdullah, I. Bello, S. Hamad & A. Bashi, "Prevalence of human leishmaniasis in Sudan: A systematic review and meta-analysis", *World J. Methodol.* **12** (2022) 305.
- [4] O. P. Singh & S. Sundar, "Visceral leishmaniasis elimination in India: progress and the road ahead", *Expert Review of Anti-infective Therapy* **20** (2022) 1381.
- [5] J. Seaman, D. Pryce, H. E. Sondorp, A. Moody, A. D. M. Bryceson & R. N. Davidson, "Epidemic Visceral Leishmaniasis in Sudan: A Randomized Trial of Aminosidine plus Sodium Stibogluconate versus Sodium Stibogluconate Alone", *The Journal of Infectious Diseases* **168** (1993) 715.
- [6] A.M. El-Hassan, M.A. Ahmed, A. A. Rahim, A.A. Satir, A. Wasfi, A. A. Kordofani, M.D. Mustafa, S. Wasfi, H. Bella & M. O. Karrar, "Visceral leishmaniasis in the Sudan: clinical and hematological features", *Annals of Saudi Medicine* **10** (1990) 51.
- [7] L.K. Makau-Barasa, D. Ochol, K.A. Yotebieng, C. B. Adera & D.K. de Souza, "Moving from control to elimination of Visceral Leishmaniasis in East Africa", *Frontiers in Tropical Diseases* (2022) 67.
- [8] A. W. Leung, *Systems of Nonlinear Partial Differential Equations: Applications to Biology and Engineering*, Springer Science & Business Media, 2013.
- [9] L. Hasan, Faeza & M. A. Abdoon, "The generalized (2+ 1) and (3+ 1)-dimensional with advanced analytical wave solutions via computational applications", *International Journal of Nonlinear Analysis and Applications* **12** (2021) 1213.
- [10] T. Roubíček, *Nonlinear Partial Differential Equations with Applications*, Birkhauser Boston, 2013.
- [11] L. Debnath, *Nonlinear Partial Differential Equations for Scientists and Engineers*, Springer Science & Business Media, 2012.
- [12] G. Jumarie, "Modified Riemann-Liouville derivative and fractional Taylor series of nondifferentiable functions further results", *Computers & Mathematics with Applications* **51** (2006) 1367.
- [13] M. A. Abdoon, F. L. Hasan & N. E. Taha, "Computational Technique to Study Analytical Solutions to the Fractional Modified KDV-Zakharov-Kuznetsov Equation", *Abstract and Applied Analysis* **2022** (2022) 2162356.
- [14] K. S. Miller & B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley 1993.
- [15] M.A. Abdoon & F.L. Hasan, "Advantages of the differential equations for solving problems in mathematical physics with symbolic computation", *Mathematical Modelling of Engineering Problems*, **9** (2022) 268.
- [16] I. M. Elmojtaba, J. Y. T. Mugisha & M. H. Hashim, "Mathematical analysis of the dynamics of visceral leishmaniasis in the Sudan", *Applied Mathematics and Computation* **217** (2010) 2567.
- [17] I Podlubny, "Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Applications", *Mathematics in Science and Engineering* **198** (1999) 340.
- [18] A. Atangana & D. Baleanu, "New fractional derivatives with nonlocal and non-singular kernel: Theory and application to heat transfer model", *Therm. Sci.* **20** (2016) 763.
- [19] M. A. Abdoon, "First integral method: a general formula for nonlinear fractional Klein-Gordon equation using advanced computing language", *American Journal of Computational Mathematics* **5** (2015) 127.
- [20] R. Saadeh, M. A. Abdoon, A. Qazza & M. Berir, "A Numerical Solution of Generalized Caputo Fractional Initial Value Problems," *Fractal and Fractional* **7** (2023) 332.
- [21] F. E. Guma, O. M. Badawy, A. G. Musa, B. O. Mohammed, M. A. Abdoon, M. Berir & S. Y. M/ Salih, "Risk factors for death among COVID-19 Patients admitted to isolation Units in Gedaref state, Eastern Sudan: a retrospective cohort study", *Journal of Survey in Fisheries Sciences* **10** (2023) 712.
- [22] S. M. Al-Zahrani, F. E. I. Elsmih, K. S. Al-Zahrani & S. Saber, "A Fractional Order SITR Model for Forecasting of Transmission of COVID-19: Sensitivity Statistical Analysis", *Malaysian Journal of Mathematical Sciences* **16** (2022) 517.
- [23] M. A. Abdoon, "Programming first integral method general formula for the solving linear and nonlinear equations", *Applied Mathematics* **6** (2015) 568.
- [24] M. A. Abdoon, R. Saadeh, M. Berir & F. E. Guma, "Analysis, modeling and simulation of a fractional-order influenza model, Alexandria Engineering Journal" **74** (2023) 231.
- [25] A. Qazza, M. Abdoon, R. Saadeh & M. Berir, "A New Scheme for Solving a Fractional Differential Equation and a Chaotic System", *European Journal of Pure And Applied Mathematics* **16** (2023) 1128.
- [26] F. Guma, O. Badawy, M. Berir & M. Abdoon, "Numerical Analysis of Fractional-Order Dynamic Dengue Disease Epidemic in Sudan", *Journal Of The Nigerian Society Of Physical Sciences* **5** (2023) 1464.
- [27] M. Elbadri, M. Abdoon, M. Berir, & D. Almutairi, "A Symmetry Chaotic Model with Fractional Derivative Order via Two Different Methods", *Symmetry* **15** (2023) 1151.
- [28] S. Al-Zahrani, F. Elsmih, K. Al-Zahrani & S. Saber, "A Fractional Order SITR Model for Forecasting of Transmission of COVID-19: Sensitivity Statistical Analysis", *Malaysian Journal Of Mathematical Sciences* **16** (2022) 517.
- [29] M. Elbadri, M. Abdoon, M. Berir & D. Almutairi, "A Numerical Solution and Comparative Study of the Symmetric Rossler Attractor with the Generalized Caputo Fractional Derivative via Two Different Method", *Mathematics* **11** (2023) 2997.