



# Mineralization, geochemical signatures, and provenance of stream sediments on the Jos Plateau, Northcentral Nigeria

Shola C. Odewumi \*

*Department of Science Laboratory Technology, University of Jos, Jos, Nigeria*

## Abstract

This study aimed at determination of mineralization, geochemical signatures and provenance of stream sediments on the Jos Plateau, northcentral Nigeria. Ten (10) stream sediments samples collected were air-dried and subjected to X-Ray Fluorescence analysis. The  $\text{SiO}_2$  value of 49.79 to 77.03 wt%,  $\text{Fe}_2\text{O}_3$  value of 1.30 to 28.84 wt%,  $\text{Al}_2\text{O}_3$  value of 5.27 to 18.25 wt%, CaO value of 0.22 to 1.88 wt%, MgO value of 0.12 to 0.92 wt%,  $\text{K}_2\text{O}$  value of 0.44 to 2.00 wt% and  $\text{Na}_2\text{O}$  value of 0.02 to 0.65 wt%. The low values of CaO (0.22-1.88 wt %) and  $\text{Na}_2\text{O}$  (0.02-0.65 wt %) could be associated with the destruction of plagioclase feldspar while low  $\text{K}_2\text{O}$  composition (0.44-2.00 wt%) could be associated with the destruction of K-feldspar during weathering. The concentration of MnO (0.06-0.12 wt%) indicates manganite enrichment, the high  $\text{TiO}_2$  value (0.63-4.25 wt%) shows rutile enrichment while  $\text{ZrO}_2$  value (0.48-18.48 wt%) indicates zircon mineralization from the underlying lithologies. The high values of pathfinder elements (Y, Zn and Nb) could be attributed to the presence of Older and Younger Granites at the upstream of the study area. Samples 4, 5, 8 and 9 have Thorium (Th) concentrations >15 ppm threshold which point to areas favorable for Th mineralization. Enrichment Factor (EF) ranges from 3.03 to 42.42 indicating moderate, significant, very high enrichment and extremely high enrichment. The Contamination Factor (CF) of 1.27 to 201.03 indicates moderate and very high contamination possibly from geogenic sources. The provenance of stream sediments is associated with felsic and mafic sources.

DOI:10.46481/jnsps.2024.2181

**Keywords:** Enrichment, Mineralization, Provenance, Weathering

## Article History :

Received: 07 July 2024

Received in revised form: 24 August 2024

Accepted for publication: 15 September 2024

Published: 29 September 2024

© 2024 The Author(s). Published by the [Nigerian Society of Physical Sciences](#) under the terms of the [Creative Commons Attribution 4.0 International license](#). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Communicated by: O. J. Abimbola


## 1. Introduction

Stream sediment materials are derived through the process of weathering of underlying rocks and transported along the drainage basin. Sediments represent the geochemical compositions of materials from the upstream drainage basin [1] and mineralogical compositions of bedrock [2]. Geochemical stud-

ies based on the chemical analysis of active stream sediments are an effective tool with several applications [3].

Major geochemical variations are controlled by source geology and provenance, as well as chemical weathering and winnowing processes, more subtle variations are a result of land use and contamination from anthropogenic activity [2]. Heavy metals are natural constituents of the earth's crust, but human activities have drastically altered their geochemical cycles [4] and heavy metals are transported via erosion into streams and rivers [5].

\*Corresponding author: Tel.: +234-803-499-1835.

Email address: [sholaodewumi@yahoo.com](mailto:sholaodewumi@yahoo.com) (Shola C. Odewumi )

Lapworth *et al.* [6] provides an overview of regional geochemical mapping using stream sediments from central and south-western Nigeria in which major geochemical variations were controlled by geology, provenance, weathering and anthropogenic activities. The elemental compositions of vegetables grown along the bank of the river Delimi at village hostel, University of Jos was reported by Odewumi *et al.* [7]. The source, contamination assessment and risk evaluation of heavy metals in the stream sediments in Ibadan SW, Nigeria was reported by Okonkwo *et al.* [8].

Several authors have used major element discrimination diagrams to discriminate the tectonic settings of stream sediments [9]. The indicator mineral chemistry has increased the contribution of stream sediments in understanding the Geology and mineral Prospectivity [10]. The present study aimed at determination of mineralization, geochemical signatures and provenance of stream sediments on Jos Plateau, northcentral Nigeria.

Lar *et al.* [11] reported on heavy metals in the urban soils, stream sediments and vegetables in Jos metropolis with implications on children's health. Several researchers have also reported on heavy metals contaminations of soils in Jos [12, 13]. However, the researches on mineralization in Jos were centred on tin mineralization for over five decades and the focus was on cassiterite, sulphide mineralization and tin mining activities.

Ogunyele & Akingboye [14] described the tin mineralization in Jos and suggested to be associated with late-stage peraluminous granites both as disseminations in veins and greisens. The zeolite mineralization within the basalts of Tim-Kwala, Jos Plateau was reported by Mangdong *et al.* [15] while Davou *et al.* [16] reported on subsurface cavity produced by tin mining activities in Jos but less attention was given to other mineralization and enrichment in stream sediments on the Jos Plateau. Therefore, there is need to determine the mineralization, contamination, enrichment and provenance of stream sediments using major and trace geochemical compositions

## 2. Geology of the study area

An outline of Geology of the area is similar to that of Jos Plateau and Geological map of the study area is shown in Figure 1. The rock units constitute part of the Basement Complex in northcentral Nigeria and include: migmatite and aplopegmatitic granite gneisses that are Precambrian to Lower Paleozoic in age while hornblende biotite porphyry is Jurassic in age of Younger Granite Complexes. Basic dyke of younger age have been identified across-cutting the Migmatite and Aplopegmatitic granite gneiss (Figure 1). The detailed geology of part of Jos-Plateau, northcentral Nigeria was reported by Odewumi *et al.* [17].

## 3. Materials and methods

### 3.1. Sample collection

The study area is located on Jos Plateau, northcentral Nigeria on latitudes 9.968° to 9.890° N and longitudes 8.884° to 8.900° E (Figure 2). A Global Positioning System (GPS)




Figure 1: Geological map of the study area [18].




Figure 2: Location map of the study area with sampling points (after Odewumi *et al.* [17]).

GARMIN 78SC Model was used to appropriately locate the sampling points on the topographical map, the longitude and latitude readings were recorded in the field note. The stream sediments were taken through systematic sampling of active streams along the drainage system in the study area. Ten (10) stream sediments were collected at the centre of the stream courses to obtain more recent and active stream sediments using the natural drainage system at an interval of about 500 m (Figure 2).

The stream sediments were collected at depth of about 30cm and labeled with appropriate sample number while the top 20-25 cm of sediment is discarded to avoid spurious high

contents of Fe and Mn in oxide coatings. Hand trowels were utilized to collect at each sampling point where the flow velocity was low. The organic materials, winnowed sediments, deposits of well-sorted gravel and sources of contamination from roads and habitation were strictly avoided.

### 3.2. Geochemical analysis

Ten (10) stream sediments samples were air-dried and subjected to X-Ray Fluorescence (XRF) for major elements and trace elements compositions of stream sediments at Nigerian Geological Survey Agency Kaduna with sample preparations, procedures and analyses [19]. The values of heavy metals from the stream sediments were compared with standards from regulatory bodies such as background reference value of UNEP [20] and permissible limit of WHO [21]. The trace elements compositions were compared with the background values of Clarke [22].

### 3.3. Statistical analysis

The results of major and trace elements compositions of stream sediments obtained from XRF analysis were subjected to ANOVA analysis using SPSS software. The spatial distribution diagrams were plotted using ArcGIS software

### 3.4. Single-element pollution indices

Single pollution indices are employed in the assessment of metal contamination because they show how concentrated an element is in a certain location in comparison to a background. As an example:

Contamination factor (CF) is a quantitative evaluation of the contaminant's level and sources. The following is how CF is assessed:  $CF = C_{m_{Sample}}/C_{m_{Background}}$

where  $C_{m_{Sample}}$  = concentration of a given metal in the sediment,  $C_{m_{Background}}$  = Background value of the metal of interest at a site [23, 24]. The following four classes were established. The contamination factor (CF) with level of contamination was classified by Hakanson [25] into four (4) groups namely: low contamination ( $CF < 1$ ), moderate contamination ( $1 \leq CF < 3$ ), considerable contamination ( $3 \leq CF < 6$ ) and very high contamination ( $CF > 6$ ).

Enrichment Factors (EF) are useful indicator of contamination status and level in the research environment [4]. As seen below, the EF computation compares each value to a certain (control sample) background level in order to identify possible sources:

$$EF = (Me/Fe)_{Sample}/(Me/Fe)_{background}, \quad (1)$$

where  $(Me/Fe)_{sample}$  = the metal to Fe ratio in the sample under study;  $(Me/Fe)_{background}$  is the natural background value of metal to Fe ratio. We used metal background values from roughly 4.5 kilometers away for this study. Iron was chosen as a normalization factor since its natural sources (1.5 percent) have a significant influence on its input [26].

The Enrichment Factor (EF) was classified according to Mmolawa et al. [27] into five (5) categories namely: deficiency to minimal enrichment ( $EF < 2$ ), moderate enrichment ( $2 \leq EF < 5$ ), significant enrichment ( $5 \leq EF < 20$ ), very high enrichment ( $20 \leq EF < 40$ ) and extremely high enrichment ( $EF \geq 40$ ).

## 4. Results

The major oxides composition of stream sediments is presented in Table 1 and trace elements composition of stream sediments is presented in Table 2. The Clarke values [22] for the indicator elements in stream sediments and range of trace elements values in present study is presented in Table 3. The  $TiO_2$  (wt%) vs  $Al_2O_3$  (wt%) binary plot of stream sediments [28] is shown in Figure 3. Table 4 shows the minimum, maximum, mean, the metal to Fe ratio in the sample under study (Me/Fe) sample, the metal to Fe ratio in the natural background value (Me/Fe) background, the background value [20] and WHO standards [21].

The  $SiO_2$  value ranges from 49.79 to 77.03 wt%,  $Fe_2O_3$  value ranges from 1.30 to 28.84 wt%,  $Al_2O_3$  value ranges from 5.27 to 18.25 wt%, CaO value varies from 0.22 to 1.88 wt%, MgO value ranges from 0.12 to 0.92 wt%,  $K_2O$  value ranges from 0.44 to 2.00 wt%,  $Na_2O$  value ranges from 0.02 to 0.65 wt%,  $TiO_2$  value ranges from 0.63 to 4.25 wt%, MnO value ranges from 0.06 to 0.12 wt% and  $ZrO_2$  value varies from 0.48 to 18.48 wt% (Table 1). Statistically, the percentage compositions of stream sediments analyzed ranges from 77.02 wt% of  $SiO_2$  to 0.02 wt% of  $Na_2O$  in location 2 and the difference was significant ( $p < 0.05$ ). The Coefficient of Variability (CV) ranges from 0.57 to 1.80 %.

V varies from 230.00 to 754.00 ppm, Cr (800.00 - 2700.00 ppm), Cu (220.00 - 650.00 ppm), Sr (5.00 - 800.00 ppm), Ba (3.25 - 67.00 ppm), Zn (233.00 - 1100.00 ppm), Ga (17.55 - 86.00 ppm), Y (80 - 800 ppm), Mo (824.00 - 5900.00 ppm), Ta (22.50 - 885.00 ppm), W (9.80 - 40.00 ppm), Hf (2.00 - 9000.00 ppm), Th (1.77 - 44.60 ppm) and Sn (199.00 - 433.50 ppm) among others (Table 2).

Trace element composition from the analysis of variance was higher (9000 ppm) for Hf as present in sample 4 and lowest (0.001 ppm) in Ce, Eu, Nd, Rb and Th across all stream sediments. Generally, sample 5 had abundance composition of trace element compared to other stream sediments with a mean composition of 547 ppm. Meanwhile, Mo had the highest abundance with a mean composition of 2596.82 ppm as evaluated across all the stream sediments.

## 5. Discussion

The low values of CaO and  $Na_2O$  may be associated with the destruction of plagioclase feldspar during weathering while low  $K_2O$  composition could be associated with the destruction of K-feldspar during weathering [29]. The high values of  $ZrO_2$  (0.48 - 18.48 wt%) and Hf (2.00 - 9000.00 ppm) indicate stream sediments that lack of clayness [30]. The high concentration of MnO (0.06 - 0.12 wt%) in the stream sediments indicates possibility of manganite enrichment in the underlying lithologies. The high  $TiO_2$  value (0.63 - 4.25 wt%) shows the possibility of rutile enrichment in underlying lithologies while  $ZrO_2$  value (0.48 - 18.48 wt%) indicates zircon enrichment in the underlying lithologies [31, 32].

The mineralizations in the stream sediments samples were derived through the weathering of minerals hosted in the under-

Table 1: Major oxide compositions (wt%) of stream sediments.

Major Oxide	1	2	3	4	5	6	7	8	9	10
SiO <sub>2</sub>	49.79a	77.02a	71.80a	59.32a	73.04a	65.19a	70.90a	62.56a	70.99a	50.37a
Fe <sub>2</sub> O <sub>3</sub>	16.64c	1.30e	3.47c	2.64e	2.18e	16.42c	4.00d	16.00b	3.34c	28.84b
Al <sub>2</sub> O <sub>3</sub>	18.25b	8.78b	14.00b	15.44c	13.00b	17.19b	8.87b	6.89c	6.53b	5.27c
CaO	0.25h	0.22h	0.40h	1.00g	1.44g	0.66f	1.88f	0.97f	1.56f	0.60g
MgO	0.44g	0.31g	0.37i	0.12i	0.56h	0.81d	0.92g	0.72g	0.87g	0.37i
K <sub>2</sub> O	1.02e	0.56f	2.00e	1.23f	1.80f	0.67e	0.54h	0.44h	0.56h	2.00d
Na <sub>2</sub> O	0.04j	0.02j	0.61g	0.43h	0.55i	0.04j	0.11j	0.20i	0.08j	0.65f
TiO <sub>2</sub>	1.15d	2.18d	1.71f	4.25d	2.60d	0.63g	2.53e	1.32d	1.80e	1.24e
MnO	0.11i	0.06i	0.06j	0.07j	0.09j	0.11i	0.12i	0.09j	0.10i	0.10k
P <sub>2</sub> O <sub>5</sub>	ND	ND	ND	ND	ND	ND	ND	ND	ND	0.36j
ZrO <sub>2</sub>	0.91f	5.16c	3.27d	18.48b	5.41c	0.12h	6.20c	1.08e	2.75d	0.48h
LSD <sub>(0.05)</sub>	0.06	0.03	0.13	0.04	0.01	0.01	0.01	0.08	0.22	0.04
CV (%)	0.58	0.97	1.63	0.58	0.57	0.32	1.80	0.70	0.65	1.68

Note: Means followed by the same letter (s) within a column are not statistically significant at 5% level of Probability; ND-Not Detected; LSD- Least Significant Difference; CV- Coefficient of Variability.

Table 2: Trace elements compositions (ppm) of stream sediments.

Trace elements	Stream sediments									
	1	2	3	4	5	6	7	8	9	10
V	700.60 <sup>lmnopq</sup>	564.00 <sup>lmnopq</sup>	366.00 <sup>nopq</sup>	754.00 <sup>lmnopq</sup>	298.00 <sup>opq</sup>	230.00 <sup>opq</sup>	330.00 <sup>opq</sup>	323.00 <sup>opq</sup>	334.00 <sup>nopq</sup>	300.00 <sup>opq</sup>
Cr	2000.00 <sup>fg hij</sup>	800.00 <sup>klmnopq</sup>	1000.00 <sup>ijklmnop</sup>	1011.00 <sup>ijklmnop</sup>	1500.00 <sup>ghijkl</sup>	1500.00 <sup>ghijkl</sup>	1400.00 <sup>hijklm</sup>	2050.00 <sup>fg hi</sup>	2700.00 <sup>def</sup>	1800.00 <sup>fg hijk</sup>
Cu	500.50 <sup>lmnopq</sup>	420.00 <sup>nopq</sup>	300.50 <sup>opq</sup>	350.00 <sup>nopq</sup>	650.00 <sup>lmnopq</sup>	405.00 <sup>nopq</sup>	220.00 <sup>opq</sup>	400.00 <sup>nopq</sup>	420.00 <sup>nopq</sup>	305.00 <sup>opq</sup>
Sr	5.00 <sup>q</sup>	800.00 <sup>klmnopq</sup>	366.00	55.00 <sup>pq</sup>	233.00 <sup>opq</sup>	12.00 <sup>q</sup>	500.00 <sup>lmnopq</sup>	54.00 <sup>pq</sup>	48.50 <sup>q</sup>	510.00 <sup>lmnopq</sup>
Ba	16.90 <sup>q</sup>	9.50 <sup>q</sup>	4.55 <sup>q</sup>	6.00 <sup>q</sup>	12.55 <sup>q</sup>	20.55 <sup>q</sup>	4.55 <sup>q</sup>	3.25 <sup>q</sup>	67.00 <sup>pq</sup>	32.50 <sup>q</sup>
Zn	501.00 <sup>lmnopq</sup>	233.00 <sup>opq</sup>	600.00 <sup>lmnopq</sup>	600.00 <sup>lmnopq</sup>	700.00 <sup>lmnopq</sup>	1100.00 <sup>ijklmno</sup>	500.00 <sup>lmnopq</sup>	540.00 <sup>lmnopq</sup>	1000.00 <sup>ijklmnop</sup>	605.00 <sup>lmnopq</sup>
Ce	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>
Pb	2200.00 <sup>fg hi</sup>	30.86 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	2500.00 <sup>defg</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	1300.00 <sup>ijklmn</sup>
Bi	12.00 <sup>q</sup>	220.00 <sup>opq</sup>	200.00 <sup>opq</sup>	205.50 <sup>opq</sup>	250.00 <sup>opq</sup>	0.001 <sup>q</sup>	270.00 <sup>opq</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	32.55 <sup>q</sup>
Ga	35.50 <sup>q</sup>	86.00 <sup>pq</sup>	49.00 <sup>q</sup>	17.55 <sup>q</sup>	22.50 <sup>q</sup>	70.00 <sup>pq</sup>	65.50 <sup>pq</sup>	22.85 <sup>q</sup>	34.00 <sup>q</sup>	40.00 <sup>q</sup>
As	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.006 <sup>q</sup>	0.001 <sup>q</sup>	0.02 <sup>q</sup>	0.001 <sup>q</sup>	0.065 <sup>q</sup>
Nd	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>
Y	100.00 <sup>pq</sup>	170.00 <sup>opq</sup>	150.00 <sup>pq</sup>	158.85 <sup>pq</sup>	200.50 <sup>opq</sup>	120.00 <sup>pq</sup>	190.00 <sup>opq</sup>	80.00 <sup>q</sup>	240.00 <sup>opq</sup>	800.00 <sup>klmnopq</sup>
Ni	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	300.00 <sup>opq</sup>	200.00 <sup>opq</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>
Rb	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>
Mo	1100.00 <sup>ijklmno</sup>	5900.00 <sup>b</sup>	2100.00 <sup>fg hi</sup>	2230.00 <sup>efg hi</sup>	3400.00 <sup>cd</sup>	2500.00 <sup>defg</sup>	4200.00 <sup>c</sup>	1300.00 <sup>ijklmn</sup>	2500.00 <sup>defg</sup>	824.00 <sup>klmnopq</sup>
Tl	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>
Cd	0.06 <sup>q</sup>	0.001 <sup>q</sup>	0.085 <sup>q</sup>	0.001 <sup>q</sup>	0.055 <sup>q</sup>	0.001 <sup>q</sup>	0.556 <sup>q</sup>	0.001 <sup>q</sup>	0.08 <sup>q</sup>	0.055 <sup>q</sup>
Ru	0.81 <sup>q</sup>	1.59 <sup>q</sup>	1.60 <sup>q</sup>	1.09 <sup>q</sup>	1.41 <sup>q</sup>	1.43 <sup>q</sup>	1.62 <sup>q</sup>	1.02 <sup>q</sup>	1.19 <sup>q</sup>	0.71 <sup>q</sup>
Eu	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>
Re	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>
Nb	700.00 <sup>lmnopq</sup>	1800.00 <sup>fg hijk</sup>	1100.00 <sup>ijklmno</sup>	2800.00 <sup>def</sup>	1800.00 <sup>ghijkl</sup>	800.80 <sup>klmnopq</sup>	2400.00 <sup>defg hi</sup>	54.00 <sup>pq</sup>	2300.00 <sup>efg hi</sup>	400.00 <sup>nopq</sup>
Ta	580.00 <sup>lmnopq</sup>	452.00 <sup>mnopq</sup>	788.00 <sup>klmnopq</sup>	685.00 <sup>lmnopq</sup>	885.00 <sup>klmnopq</sup>	435.00 <sup>nopq</sup>	452.00 <sup>mnopq</sup>	22.50 <sup>q</sup>	588.90 <sup>lmnopq</sup>	340.00 <sup>nopq</sup>
W	12.00 <sup>q</sup>	9.80 <sup>q</sup>	40.00 <sup>q</sup>	33.50 <sup>q</sup>	38.00 <sup>q</sup>	20.50 <sup>q</sup>	31.00 <sup>q</sup>	40.00 <sup>q</sup>	22.55 <sup>q</sup>	13.50 <sup>q</sup>
Hf	3261.00 <sup>cde</sup>	900.00 <sup>klmnopq</sup>	600.50 <sup>lmnopq</sup>	9000.00 <sup>a</sup>	8700.00 <sup>a</sup>	38.80 <sup>q</sup>	22.54 <sup>q</sup>	2.00 <sup>q</sup>	588.00 <sup>lmnopq</sup>	12.00 <sup>q</sup>
Yb	0.87 <sup>q</sup>	0.001 <sup>q</sup>	0.08 <sup>q</sup>	0.054 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.022 <sup>q</sup>	0.001 <sup>q</sup>
Se	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.055 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.08 <sup>q</sup>	0.001 <sup>q</sup>	0.062 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>
U	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.40 <sup>q</sup>	2.55 <sup>q</sup>	0.32 <sup>q</sup>	1.02 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>
Th	2.88 <sup>q</sup>	2.70 <sup>q</sup>	12.00 <sup>q</sup>	44.60 <sup>q</sup>	43.00 <sup>q</sup>	1.77 <sup>q</sup>	3.00 <sup>q</sup>	22.50 <sup>q</sup>	32.00 <sup>q</sup>	0.001 <sup>q</sup>
Sb	0.55 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.76 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.550 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.075 <sup>q</sup>
Ge	1.05 <sup>q</sup>	4.00 <sup>q</sup>	3.55 <sup>q</sup>	6.22 <sup>q</sup>	7.70 <sup>q</sup>	52.00 <sup>pq</sup>	11.00 <sup>q</sup>	30.50 <sup>q</sup>	28.50 <sup>q</sup>	12.00 <sup>q</sup>
Sn	355.00 <sup>mnopq</sup>	275.00 <sup>opq</sup>	857.00 <sup>klmnopq</sup>	199.00 <sup>opq</sup>	221.00 <sup>opq</sup>	433.50 <sup>lmnopq</sup>	275.00 <sup>opq</sup>	654.00 <sup>lmnopq</sup>	231.00 <sup>opq</sup>	422.00 <sup>opq</sup>
Pd	0.42 <sup>q</sup>	0.48 <sup>q</sup>	0.78 <sup>q</sup>	0.55 <sup>q</sup>	0.88 <sup>q</sup>	1.10 <sup>q</sup>	0.52 <sup>q</sup>	0.001 <sup>q</sup>	1.39 <sup>q</sup>	0.25 <sup>q</sup>
La	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.005 <sup>q</sup>	0.007 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.005 <sup>q</sup>
Co	20.00 <sup>q</sup>	7.50 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	0.001 <sup>q</sup>	120.00 <sup>pq</sup>	0.001 <sup>q</sup>	17.00 <sup>q</sup>	0.566 <sup>q</sup>	310.00 <sup>opq</sup>
LSD <sub>(0.05)</sub>	863									
SE±	363.72									
CV(%)	109.82									

Means that follow the same letter (s) within a column are not statistically significant at 5% level of probability. CV: Coefficient of Variability; SE± : Standard Error of Means; LSD<sub>(0.05)</sub>: Least Significant Difference at 5% level of probability.

lying granitic rocks upstream of the study area that have been dispersed along the stream. This is similar to the report of Lapworth *et al.* [6] on geochemical mapping using stream sedi-

ments in west-central Nigeria where high stream sediment Zr concentrations (mean >0.2%), from proximal zircons derived from weathering of basement rocks.

Table 3: Clarke [22] values for the indicator elements in stream sediments and range of elements' values in present study.

Element	Clarke (ppm)	Range of elements
Zr	200	888.00-136,808.00
Nb	10	54.00 -2800.00
Zn	25	233.00 - 1100.00
Y	18	80 - 800 ppm

The mineralizations in the study area are similar to the report of Ayodele [5] on Stream Sediment of Ara, Epe and Ijero Area, South Western Nigeria that was mineralized in phosphate minerals such as monazite, turquoise, or apatite; ferruginous minerals which could be hematite and manganese-bearing minerals such as manganite or mica rich manganese. These mineralizations in Ara, Epe and Ijero Area are hosted by the pegmatites that intrude the country rocks in the northwestern part of the area.

The  $Fe_2O_{3T}$  value ranges from 1.30 to 28.84 wt% (Table 1). Some stream sediments samples like sample 1 (16.64 wt %), sample 6 (16.42 wt%), sample 8 (16.00 wt%) and sample 10 (28.84 wt%) have  $Fe_2O_{3T} > 15$  % and can be classified as Fe-rich sediments which could have been mineralized by the underlying Fe-rich mineralized rocks and veins. Low grade ore is a term applied to iron-rich rocks with cut-off grades in the range of 15-25%  $Fe_2O_{3T}$ . According to Levinson [33], hematite mineralization can be associated with circulating hydrothermal fluids during late stages of cooling of molten magma. This is similar to the report of Ayodele [5] on stream sediments from Ara, Epe and Ijero Area, South Western Nigeria where ferruginous minerals were reported.

The Sn value ranges from 199.00 to 654.00 ppm and could be linked to cassiterite mineralization possibly from Younger Granitic rocks upstream of the study area. The high concentrations of pathfinder elements such as Y (80.00 - 800.00 ppm), Zn (233.00 -1100.00ppm) and Nb (54.00 -2800.00 ppm) were obtained from the study area (Table 2) and could be attributed to the presence of Older and Younger granitic rock at the upstream of the study area. This is similar to the report of Mira *et al.* [2] on Stream Sediments Geochemical Exploration of Wadi El Reddah area, Northeastern Desert, Egypt where high contents of pathfinder elements (REE, Y, Zn, Nb and As) were discovered and attributed to the presence of alkali feldspar granite at Gabal Gattar area upstream of Wadi El Reddah in Egypt.

The Th value ranges from <0.001 to 44.60 ppm (Table 2) and Samples 4, 5, 8 and 9 have Th values greater than threshold value of 15 ppm. This point to some of the areas favorable for Th mineralization and can be linked to remobilization and dispersion of Thorite within the stream sediments while U value ranges from <0.001 to 2.55 ppm (Table 2) indicating lack of U mineralization in the area [22]. This is in contrast to the report of Mira *et al.* [2] on Stream Sediments from Egypt where >5 ppm U and >15 ppm Th as threshold were reported and pointed to areas favorable for U mineralization that was linked to deformation-induced radioelement remobilization and subsequent precipitation in veins forms.

Table 3 shows anomalous concentrations of pathfinder elements compared with the published background values of [20]. Higher value of Zr (0.48 - 18.48 wt%) in the stream sediments of the study area than the Zr value for indicator element of 200ppm [22] suggests zircon mineralization. Higher values of Y (80 - 800 ppm) in the stream sediments of the study area than the Y indicator element value of 18 ppm [22] point to xenotime mineralization in the area while higher values of Nb (54.00 - 2800.00ppm) in the stream sediment than Nb indicator value of 10ppm [22] point to columbite mineralization (Table 3). The stream sediments have high values of Ta ranging from 22.50 to 885.00 ppm (Table 2) which point to enrichment of tantalite in the study area.

The study area is enriched in resisting minerals such as zircon, rutile, xenotime, thorite, cassiterite, and columbite-tantalite that could be attributed to the underlying mineralization hosted within the Older and Younger Granitic rocks in the area that were dispersed along the drainage basin. This is similar to the report of Mira *et al.* [2] on Stream Sediments from Egypt where zircon, xenotime, thorite, and columbite-tantalite mineralizations were reported and hosted by alkali feldspar granite

The research of Lapworth *et al.* [6] on geochemical mapping of stream sediments in west-central Nigeria identified placer deposits of potential economic importance for Au, REE, Ta, Nb, U and Pt, as well as other primary metal deposits. The values As (<0.001 -0.065 ppm) and Cr (800 – 2000 ppm) obtained in the study area (Table 2) were possibly derived from Jurassic Younger Granites of Jos Plateau. This is similar to the report of Lapworth *et al.* [6] where higher As (>2 ppm) and Cr (>70 ppm) were associated with Mesozoic and younger coastal stream sediments of west-central Nigeria.

Enrichment Factor (EF) ranges from 3.03 to 42.42, this indicates moderate enrichment ( $2 \leq EF < 5$ ), significant enrichment ( $5 \leq EF < 20$ ), very high enrichment ( $20 \leq EF < 40$ ) and extremely high enrichment ( $EF \geq 40$ ) according to enrichment categories of Mmolawa *et al.* [27]. The higher average values of Cu (397.10 ppm) than the WHO [21] standard (25 ppm) indicates contamination. Similarly, higher average values of Zn, Ni and Cr than the WHO standards [21] attest to contamination in the study area (Table 4). Contamination Factor (CF) ranges from 1.27 to 201.03 which indicate moderate contamination ( $1 \leq CF < 3$ ) and very high contamination ( $CF > 6$ ) according to contamination level of Hakanson [25]. The contamination could be attributed to geogenic sources

The presence and distribution of heavy metals in soils are influenced mainly by the parent material, the chemical and physical soil properties, the metal speciation, and the climatic conditions. The mineral content of the parent material is one of the most important factors for the amount of trace elements in soils, irrespective of classification or the amount of weathering [34]. This was observed in the present study where heavy metals in the stream sediments were suggested to be derived from geogenic sources.

Anthropogenic sources of contaminations from industrial, mining and domestic activities [35, 36] could not be ascertained in the study area along the stream sediments since an-

Table 4: Minimum, maximum, mean, the metal to Fe ratio in the sample under study  $(Me/Fe)_{(s)}$  the metal to Fe ratio in the natural background value  $(Me/Fe)_{(brd)}$ , Enrichment Factor (EF), Contamination Factor (CF), background value [20] and WHO standard [21].

Elements	Min.	Max.	Mean	$Me/Fe_{(s)}$	$/Me/Fe_{(brd)}$	ER	CF	Background value	WHO
Cu	220.0	650.00	397.10	41.89	6.91	6.06	198.55	2	25
Pb	<0.001	2500	603.09	63.61	6.9	9.09	201.03	3	----
Zn	233	1100	637.90	67.19	3.17	21.21	91.13	7	123
Ni	0.001	300	50.00	5.27	1.74	3.03	50	1	20
Co	0.001	310	47.51	5.01	0.82	6.06	23.76	2	---
Sr	5.00	800.00	258.35	27.25	3.00	9.09	86.12	3	----
V	230.00	754.00	419.90	44.29	1.62	27.27	46.66	9	----
Cr	800	2700	1,511.10	159.40	5.26	30.30	151.11	10	25
Ba	3.25	67.00	17.74	1.87	0.044	42.42	1.27	14	---
Fe (wt%)	1.30	28.84	9.48					0.33	




Figure 3:  $TiO_2$  (wt%) vs  $Al_2O_3$  (wt%) binary plot of stream sediments [28].

thropogenic sources are usually referred to as ‘‘Punctual’’ contamination with an identified source often close to the contaminated soil [37]. This is in contrast to the reports of Odewumi *et al.* [38] and Momoh *et al.* [39] on soils and vegetables from Jos where anthropogenic and/or geogenic sources of contamination were suggested. Okonkwo *et al.* [8] reported on stream of Rivers around Olode Area SW, Nigeria where anthropogenic sources of contamination from mining activities which had negative impact on the study area have been indicated.

On the  $TiO_2$  (wt%) versus  $Al_2O_3$  (wt%) binary plot [28] as shown in Figure 3, the stream sediments plot in the basalt, basalt+granite/rhyolite, granite/rhyolite+basalt and granite/rhyolite fields. This indicates that the provenance could be from felsic and mafic sources.

The stream sediments have low percentages of MgO, CaO,  $K_2O$  and  $Na_2O$  and may be attributed to intense weathering and alteration of K-feldspar, plagioclase feldspar and unstable mafic minerals [40, 41]. The relatively high values of the ratio of  $K_2O/Na_2O$  could be attributed to the presence of albitic plagioclase, K-feldspar, mica and illite [42] which is a major characteristic of felsic sources. The low value of  $Na_2O$  (0.02 to 0.65 wt%) in the stream sediments point to the relatively small amount of Na-rich plagioclase in the protoliths of the stream sediments [43].

The geochemical signature in stream sediments is strongly




Figure 4: Spatial distribution plot of  $ZrO_2$  (wt%).




Figure 5: Spatial distribution plot of  $MnO$  (wt%).

influenced by the geological setting and local lithologies in




Figure 6: Spatial distribution diagram of  $\text{TiO}_2$  (wt %).Figure 7: Spatial distribution diagram of  $\text{Fe}_2\text{O}_3$  (wt %).

Figure 8: Spatial distribution diagram of Sn (ppm).




Figure 9: Spatial distribution diagram of Y (ppm).

Brazil [44]. This is similar to the present study where the geochemical signature of stream sediments is influenced by underlying mineralization and lithologies. The positive correlation between  $\text{Al}_2\text{O}_3$  and  $\text{Fe}_2\text{O}_3$ ,  $\text{MnO}$ , and  $\text{MgO}$ , indicates multiple sources for sediment and these oxides are associated with clay minerals of Eastern Ghats Mobile Belt in India [45]. The present findings also show multiples sources of stream sediments on the Jos Plateau which could be attributed to felsic and mafic sources.

The discrimination function diagram suggests a felsic igneous as well as quartzo-sedimentary provenance for stream sediments in India whereas the ratio of  $\text{Al}_2\text{O}_3/\text{TiO}_2$  and concentration of V-Ni-Th suggest a felsic igneous source rock [45]. Similarly, the relatively high values of the ratio of  $\text{K}_2\text{O}/\text{Na}_2\text{O}$  obtained in the present study could be attributed to felsic

sources. The low concentrations of  $\text{MgO}$  (mean 0.03%),  $\text{Na}_2\text{O}$  (mean 0.008%), and  $\text{K}_2\text{O}$  (mean 0.04%) were reported from sandstone sediments of the Ilaro formation, Dahomey Basin, Southwestern Nigeria and suggested chemical destruction in an oxidizing environment [46]. In the present study, the low values of  $\text{CaO}$  (0.22-1.88 wt %) and  $\text{Na}_2\text{O}$  (0.02-0.65 wt %) could be associated with the destruction of plagioclase feldspar while low  $\text{K}_2\text{O}$  composition (0.44-2.00 wt%) could be associated with the destruction of K-feldspar during weathering

According to Seydi *et al.* [47], the stream sediments in East Iran were reported to be favorable for copper and gold mineralization using concentration number. This is in contrast to the present study where stream sediments on the Jos Plateau were found to be favorable for Mn, Zr, Y, Nb, Th, Sn, Ta and




Figure 10: Spatial distribution diagram of Th (ppm).




Figure 12: Spatial distribution diagram of Ta (ppm).




Figure 11: Spatial distribution diagram of Nb (ppm).

Ti mineralization. Osei *et al.* [48] suggested a stronger positive correlation of gold with Iron, Chromium and Vanadium using multivariate analysis of the geochemical data using Pearson product-moment of correlation and indicated gold mineralization in Ghana. The present study also show positive correlation of major oxides as presented in Table 1.

The enrichments of heavy metals of sediments in Mexico were identified through the accumulation and indicated that the contamination of Cd and Pb is mainly of anthropogenic origin of stream sediments in Mexico [49] while high values of Fe (>0.4 wt%), and Mn (>2000 ppm) were reported by Kelley *et al.* [50] on stream-sediment from eastern Alaska, USA and suggested geogenic sources This is in contrast to the cur-

rent findings where contamination of heavy metals in the stream sediment was predominantly from geogenic sources

According to Ajaha *et al.* [51], the concentration of the metals in the soil of Ebonyl State Nigeria such as Zn (7.28-11.33), Cu (3.40 - 4.64), Fe (803.04 - 735.47) and Cd (1.14) were within FEPA and FAO/WHO limits. This is in contrast to the present findings where the value of Zn, Cu, Fe and Cd in the stream sediments exceeded the FAO/WHO limits. According to Olutona [52], the As and Sr, values were below the background values for typical soil while the present study shows that the values of As and Sr exceeded the background values

The spatial distribution plot of  $ZrO_2$  is shown in Figure 4, the spatial distribution plot of  $MnO$  is shown in Figure 5, the spatial distribution plot of  $Fe_2O_3$  is shown in Figure 6 while Figure 7 shows the spatial distribution diagram of  $TiO_2$  which are favorable for zircon, manganite, hematite and rutile mineralizations respectively. The spatial distribution plot of Sn is shown in Figure 8, the spatial distribution plot of Y is shown in Figure 9, the spatial distribution plot of Th is shown in Figure 10, the spatial distribution diagram of Nb is shown in Figure 11 while Figure 12 shows the spatial distribution diagram of Ta which are favorable for cassiterite, xenotime, thorite, columbite and tantalite mineralizations, respectively.

## 6. Conclusion

The stream sediment is enriched in resisting minerals such as zircon, rutile, xenotime, thorite, cassiterite, and columbite-tantalite which could be attributed to the mineralization hosted in the underlying rocks. These mineralizations in the stream sediments were derived through the weathering of minerals hosted in the underlying Older and Younger Granitic rocks upstream of the study area which have been dispersed along the

drainage basin. The provenance of stream sediments from the study area could be associated with felsic and mafic sources. The relatively high values of the ratio of  $K_2O/Na_2O$  could be attributed to felsic sources. The Enrichment Factor indicates moderate enrichment, significant enrichment, very high enrichment and extremely high enrichment. The higher average values of Cu, Zn, Ni and Cr in the stream sediments samples than the WHO standards attest to contamination in the study area. The Contamination Factor (CF) indicates moderate contamination and very high contamination possibly from geogenic sources

## Acknowledgment

I write to acknowledge Mr Afizu Mamudu of Physics Department, University of Jos for plotting the spatial distribution diagrams and Mr Ife of the Department of Plant Science and Biotechnology, University of Jos for statistical analysis.

## References

- [1] M. E. Doherty, K. Arndt, Z. Chang, K. Kelley & O. Lavin, "Stream sediments geochemistry in mineral exploration: a review of fine fraction, clay-fraction, bulk leach gold, heavy mineral concentrate and indicator mineral chemistry", *Geochemistry: Exploration, Environment, Analysis* **23** (2023) 1. <https://doi.org/10.1144/geochem2022-039>.
- [2] H. Mira, H. Hussein, S. Tawfik & N. Abed, "Stream sediments geochemical exploration in Wadi El Reddah area, Northeastern desert, Egypt", *Mediterranean Journal of Chemistry* **10** (2020) 809. <http://dx.doi.org/10.13171/mjc10802011021539st>.
- [3] E. C. Grunsky, L. J. Drew & D. M. Sutphin, "Process recognition in multi-element soil and stream-sediment geochemical data", *Applied Geochemistry* **24** (2009) 1602. <http://dx.doi.org/10.1016/j.apgeochem.2009.04.024>
- [4] H. Feng, X. Han & W. G. Zhang, "A preliminary study of heavy metal contamination in Yangtze River Intertidal Zone Due to urbanization", *Marine Pollution Bulletin* **49** (2004) 910. <https://doi.org/10.1016/j.marpolbul.2004.06.014>.
- [5] O. S. Ayodele, "Stream sediment geochemical survey of Ara, Epe and Ijoro area, South Western Nigeria", *International Journal of Science and Technology* **1** (2011) 269. <https://api.semanticscholar.org/CorpusID:14216297>.
- [6] D. J. Lapworth, K. V. Knights & R. M. Key, "Geochemical mapping using stream sediments in west-central Nigeria: Implications for environmental studies and mineral exploration in West Africa", *Applied Geochemistry* **27** (2012) 1035. <https://doi.org/10.1016/j.apgeochem.2012.02.023>.
- [7] S. C. Odewumi, I. D. Yohana, J. A. Bulus & I. Ogbe, "Geochemical appraisals of elemental compositions of stream sediments and some vegetables at Village Hostel University of Jos", *Nigerian Annals of Pure and Applied Science* **3** (2020) 77. <https://doi.org/10.46912/napas.150>.
- [8] S. I. Okonkwo, S. O. Idakwo, M. S. Kolawole, O. Faloye & A. A. Elueze, "Source, contamination assessment and risk evaluation of heavy metals in the stream sediments of rivers around Olode area SW, Nigeria", *Journal of Environmental & Earth Sciences* **5** (2023) 65. <https://doi.org/10.30564/jees.v5i1.5060>.
- [9] M. R. Bhatia, "Plate tectonics and geochemical composition of sandstone", *Journal of Geology* **91** (1983) 611. <http://dx.doi.org/10.1086/628815>.
- [10] J. S. Armstrong-Altrin, Y. I. Lee, S. P. Verma & S. Ramasamy, "Geochemistry of sandstones from the upper Miocene kudankulam Formation, southern India: Implication for provenance, weathering and tectonic setting", *Journal of Sedimentary Research* **74** (2004) 285. <http://dx.doi.org/10.1306/082803740285>.
- [11] U. Lar, E. Shettima & H. Dibal, "Heavy metals in the urban soils and vegetables in Jos metropolis, Nigeria: implications on children's health", *American Journal of Environmental Protection. Special Issue: Integrating Earth Materials, Diet, Water and Human Health* **3** (2014) 70. <http://doi:10.11648/j.ajep.s.2014030602.20>.
- [12] S. C. Odewumi, C. I. Dihis & A. A. Idris, "Geochemical and mineralogical compositions of geophagic clays from terminus and new market, Jos, Plateau State, Nigeria", *Fuwakari Journal of trends in Science and Technology* **8** (2023) 255. [https://www.researchgate.net/publication/382365507\\_GEOCHEMICAL\\_AND\\_MINERALOGICAL\\_COMPOSITIONS\\_OF\\_GEOPHAGIC\\_CLAYS\\_FROM\\_TERMINUS\\_AND\\_NEW\\_MARKET JOS PLATEAU STATE NIGERIA](https://www.researchgate.net/publication/382365507_GEOCHEMICAL_AND_MINERALOGICAL_COMPOSITIONS_OF_GEOPHAGIC_CLAYS_FROM_TERMINUS_AND_NEW_MARKET JOS PLATEAU STATE NIGERIA).
- [13] S. C. Odewumi, M. A. Onimisi, M. O. Adeoye, A. N. Change & B. T. Omoyajowo, "Palaeoweathering, provenance and hydrothermal alteration characteristics of Nahuta clay, Jos-Plateau, Northcentral Nigeria", *Journal of Environmental and Earth Sciences* **6** (2024) 164. <https://doi.org/10.30564/jees.v6i2.6286>.
- [14] A. C. Ogunyele & A. S. Akingboye, "Tin mineralisation in Nigeria: a review", *Environmental and Earth Sciences Research Journal* **5** (2018) 15. <https://doi.org/10.18280/eesrj.050103>.
- [15] C. L. Mangdong, K. C. Aluwong, M. P. Goyit & D. A. Bala, "Zeolite mineralization prospect within the Basalts of Tim-Kwalla, Jos Plateau North Central Nigeria", *American Journal of Engineering Research* **6** (2017) 64. <https://api.semanticscholar.org/CorpusID:210612736>.
- [16] H. D. Davou, S. C. Odewumi & E. S. Akanbi, "Subsurface cavity detection using 2d electrical resistivity tomography at some mining sites In Jos, Northcentral Nigeria", *FUPRE Journal of Scientific and Industrial Research* **8** (2024) 208. <https://journal.fupre.edu.ng/index.php/fjsir/article/view/284>.
- [17] S. C. Odewumi, "Geological settings and geochemistry of younger granitic rocks from Kuba area, Ropp Complex, northcentral Nigeria", *FUPRE Journal of Scientific and Industrial Research* **4** (2020) 9. <https://journal.fupre.edu.ng/index.php/fjsir/article/download/98/75/>.
- [18] NGSA, "Geological and mineral resources map album", Nigerian Geological Survey Agency, Ministry of Solid Minerals Development. [Online]. [https://miningdecision.minesandsteel.gov.ng/ngsa\\_metadata/srv/search?serviceName=Nigerian%20Geological%20Survey%20Agency](https://miningdecision.minesandsteel.gov.ng/ngsa_metadata/srv/search?serviceName=Nigerian%20Geological%20Survey%20Agency).
- [19] S. C. Odewumi, "Geochemical characteristics of Balanga Limestone, North-Eastern Nigeria: Implications for Post-Depositional Alteration", *African Journal of Natural Sciences* **23** (2020) 83. [https://www.researchgate.net/publication/383213368\\_GEOCHEMICAL\\_CHARACTERISTICS\\_OF\\_BALANGA\\_LIMESTONE\\_NORTH-EASTERN\\_NIGERIA\\_IMPLICATIONS\\_FOR\\_POST-DEPOSITIONAL ALTERATION](https://www.researchgate.net/publication/383213368_GEOCHEMICAL_CHARACTERISTICS_OF_BALANGA_LIMESTONE_NORTH-EASTERN_NIGERIA_IMPLICATIONS_FOR_POST-DEPOSITIONAL ALTERATION).
- [20] UNEP, "Environmental risks and challenges of anthropogenic metals flows and cycles", A Report of the Working Group on the Global Metal Flows to the International Resource Panel, 2013. [Online]. <https://www.resourcepanel.org/reports/environmental-risks-and-challenges-anthropogenic-metals-flows-and-cycles>
- [21] WHO, "Global health observatory data repository", Burden of disease. Lead attributable DALYs. [Online]. <http://apps.who.int/gho/data/node.home>.
- [22] F. W. Clarke, *The relative abundance of the chemical elements*, Bulletin of the Philosophical Society of Washington, 1889, pp. 135 - 143. [https://books.google.com.ng/books/about/The\\_Relative\\_Abundance\\_of\\_the\\_Chemical\\_E.html?id=JJEtHQACAAM&redir\\_esc=y](https://books.google.com.ng/books/about/The_Relative_Abundance_of_the_Chemical_E.html?id=JJEtHQACAAM&redir_esc=y).
- [23] J. P. Brady, G. A. Ayoko & W. N. Martens, "Development of a hybrid pollution index for heavy metals in marine and estuarine sediments", *Environmental Monitoring and Assessment* **187** (2015) 306. <https://doi.org/10.1007/s10661-015-4563-x>.
- [24] G. Qingjie, D. Jun & X. Yunchuan, "Calculating pollution indices by heavy metals in ecological geochemistry assessment and a case study in parks of Beijing", *Journal of China University of Geosciences* **19** (2008) 230. [https://doi.org/10.1016/S1002-0705\(08\)60042-4](https://doi.org/10.1016/S1002-0705(08)60042-4).
- [25] L. Hakanson, "An ecological risk index for aquatic pollution control a sedimentological approaches". *Water Research* **14** (1980) 975. [https://doi.org/10.1016/0043-1354\(80\)90143-8](https://doi.org/10.1016/0043-1354(80)90143-8).
- [26] V. K. Tippie, "An environmental characterization of Chesapeake Bay and a frame work for action", in *The estuary as a filter*, Academic Press, New York, 1984, pp. 467-487. <https://doi.org/10.1016/B978-0-12-405070-9.50028-1>.

- [27] K. Mmolawa, A. Likuku & G. Gaboutloeloe, "Assessment of heavy metal pollution in soils along roadside areas in Botswana", *African Journal of Environmental Science and Technology* **5** (2011) 186. <https://www.ajol.info/index.php/ajest/article/view/71926>.
- [28] G. E. Ekosse, "Provenance of the Kgwakgwe kaolin deposit in south eastern Botswana and its possible utilization", *Applied Clay Science* **20** (2001) 137. [https://doi.org/10.1016/S0169-1317\(01\)00064-3](https://doi.org/10.1016/S0169-1317(01)00064-3).
- [29] S.C. Odewumi, "Mineralogy and geochemistry of geophagic clays from Share Area, Northern Bida sedimentary basin, Nigeria", *J Geol Geosci* **2** (2013) 108. <http://dx.doi.org/10.4172/2329-6755.1000108>.
- [30] N. N. Chombong, C. E. Suh & D. C. Ilouga, "New detrital zircon U-Pb ages from BIF related metasediments in the Ntem Complex (Congo craton) of southern Cameroon, West Africa", *Natural Science Journal* **5** (2013) 835. <http://dx.doi.org/10.4236/ns.2013.57101>.
- [31] S. C. Odewumi & V.O. Olarewaju, "Petrogenesis and geotectonic settings of the granitic rocks of Idofin-osi-eruku Area, Southwestern Nigeria using trace element and rare earth element geochemistry", *Journal of Geology & Geosciences* **2** (2013) 1. <http://dx.doi.org/10.4172/2329-6755.1000109>.
- [32] S. C. Odewumi, A. A. Aminu, A. Momoh & J. A. Bulus, "Environmental impact of mining and pedogeochemistry of Agunjin area, southwestern Nigeria", *International Journal of Earth Sciences and Engineering* **8** (2015) 558. [https://www.researchgate.net/publication/286469485\\_Environmental\\_impact\\_of\\_mining\\_and\\_pedogeochemistry\\_of\\_Agunjin\\_area\\_southwestern\\_Nigeria](https://www.researchgate.net/publication/286469485_Environmental_impact_of_mining_and_pedogeochemistry_of_Agunjin_area_southwestern_Nigeria).
- [33] A. A. Levinson, *Introduction to exploration geochemistry*, Applied Publishing Ltd., Wilmette, 1980, pp. 89-91. [https://books.google.com.ng/books/about/Introduction\\_to\\_Exploration\\_Geochemistry.html?id=g4A35GK7nOIC&redir\\_esc=y](https://books.google.com.ng/books/about/Introduction_to_Exploration_Geochemistry.html?id=g4A35GK7nOIC&redir_esc=y).
- [34] B. J. Alloway, *Heavy metal in soils*, John Wiley and Sons Incorporated, London, 1996, pp. 149-159. <https://link.springer.com/book/10.1007/978-94-007-4470-7>.
- [35] A. Migdal, E. Drag-Kozak, J. Kania-Gierdziewicz & L. Migdal, "Correlation between age and levels of heavy metals in mares mammary gland secretions", *Anim. Sci. Genet.* **19** (2023) 95. <http://dx.doi.org/10.5604/01.3001.0053.5979>.
- [36] M. Sudhakaran, D. Ramamoorthy, V. Savitha & S. Balamurugan, "Assessment of trace elements and its influence on physico-chemical and biological properties in coastal agroecosystem soil, Puducherry region", *Geology, Ecology and Landscapes* **2** (2018) 169. <https://doi.org/10.1080/24749508.2018.1452475>.
- [37] K. G. Serelis, I. G. Kafkala, K. Parpodis & S. Lazaris, "Anthropogenic and Geogenic contamination due to heavy metals in the vast area of Vari, Attica", *Bulletin of the Geological Society of Greece* **43** (2010) 2390. <https://doi.org/10.12681/bgsg.11639>.
- [38] S. C. Odewumi, O. Q. Ajegba, J. A. Bulus & I. Ogbe, "Assessment of heavy metal contaminations of soils from dumpsites in Jos Metropolis, Plateau State, Nigeria", *FULAFIA Journal of Science and Technology* **6** (2020) 37. <https://lafiascijournals.org.ng/index.php/fjst/article/view/182>.
- [39] A. Momoh, E. P. Rotji, S. C. Odewumi, M. Opuwari, O. J. Ojo & A. Olorunyomi, "Preliminary investigation of trace elements in Acid Mine Drainage from Odagbo Coal Mine, northcentral, Nigeria", *Journal of Environment and Earth Science* **7** (2017) 90. <https://api.semanticscholar.org/CorpusID:212611971>.
- [40] W. Getaneh, "Geochemistry provenance and depositional tectonic setting of the Adigrat Sandstone, northern Ethiopia", *Journal of African Earth Sciences* **35** (2002) 185. [https://doi.org/10.1016/S0899-5362\(02\)00126-4](https://doi.org/10.1016/S0899-5362(02)00126-4).
- [41] K. J. Zhang, "Secular geochemical variations of the Lower Cretaceous siliciclastic rocks from central Tibet (China) indicating a tectonic transition from continental collision to back arc rifting", *Earth and Planetary Sci. Lett.* **229** (2004) 73. <https://doi.org/10.1016/j.epsl.2004.10.030>.
- [42] B. N. Nath, H. Kunzendorf & W. L. Pluger, "Influence of provenance, weathering and sedimentary processes on the elemental ratio of the fine grained fraction of the bed load sediments from the Veinbanad Lake and the adjoining Continental shelf, southwest coast of India." *Journal of Sedimentary Research* **70** (2000) 1081. <https://doi.org/10.1306/100899701081>.
- [43] O. J. Ojo, S. A. Adepoju & D.A. Adedoyin, "Combined petrographic and geochemical studies of maastrichtian patti formation sandstone, Bida basin, Nigeria: Implications for Provenance and Compositional Maturity", *Ilorin Journal of Science* **2** (2015) 163. <https://doi.org/10.54908/ilj.2015.02.01.012>.
- [44] G. N. Salomao, R. Dall'Agnol, P. K. Sahoo, R. S. Angelica, C. A. de Medeiros Filho, J. Ferreira Junior, M. S. da Silva, P. W. Martins e Souza Filho, W. R. Nascimento Junior, M. F. da Costa, L. R. Guimaraes Guilherme & J. O. de Siqueira, "Geochemical mapping in stream sediments of the Carajas mineral province: Background values for the Itacaiunas River watershed, Brazil", *Applied Geochemistry* **118** (2020) 104608. <http://dx.doi.org/10.1016/j.apgeochem.2020.104608>.
- [45] D. K. Patel & S. Nath, "Geochemistry of stream sediments from eastern ghats mobile belt, India: implications on sediment type, maturity, source-area weathering and provenance", *Journal of Geological Society of India* **99** (2023) 1361. <https://doi.org/10.1007/s12594-023-2483-x>.
- [46] A. Y. Jimoh, M. B. Saadu, A. A. Adetoro, J. Ajadi, T. Issa & U. Issa, "Sedimentological and geochemical evaluation of sandstones of the Ilaro formation, Dahomey basin, Southwestern Nigeria : Insights into paleoenvironments, provenance, and tectonic settings", *Journal of the Nigerian Society of Physical Sciences* **6** (2024) 1726. <https://doi.org/10.46481/jnsps.2024.1726>.
- [47] A. Seydi, M. Abedi, A. Bahroudi & H. Ferdowsi, "Geochemical prospectivity of Cu-mineralization through concentration-number fractal modeling and Prediction-Area plot: a case study in east Iran", *International Journal Mining & Geo-Engineering* **57** (2023) 159. [https://ijmge.ut.ac.ir/article\\_90236.html](https://ijmge.ut.ac.ir/article_90236.html).
- [48] K. P. Osei, M. Affam, C. Kusi-Manu & D. T. Kwapong, "Multivariate statistical evaluation of geochemical data from quartz vein-associated gold mineralisation at the badukrom prospect within the Tarkwaian Base", *Ghana Mining Journal* **21** (2021) 22. <https://doi.org/10.4314/gm.v21i1.3>.
- [49] G. Navarrete-Rodríguez, M. Castañeda-Chávez & F. Lango-Reynoso, "Geoaccumulation of heavy metals in sediment of the Fluvial-Lagoon-Deltaic system of the Palizada river, Campeche, Mexico", *Int. J. Environ. Res. Public Health* **17** (2020) 969. <https://doi.org/10.3390/ijerph17030969>.
- [50] K. D. Kelley, G. E. Graham, K. Pfaff, H. A. Lowers & A. E. Koenig, "Indicator mineral analyses of stream-sediment samples using automated mineralogy and mineral chemistry: Applicability to exploration in covered terranes in eastern Alaska, USA", *Ore Geology Reviews* **148** (2022) 105021. <https://doi.org/10.1016/j.oregeorev.2022.105021>.
- [51] D. N. Ajaha, E. Agboeze, J. N. Ihediohab, E. Chukwudi-Madua & C. C. Chime, "Levels of Zinc (Zn), Copper (Cu), Iron (Fe), and Cadmium (Cd) in soil, rice stalk, and oryza sativa grain in Ishiagu rice field, Ebonyi State, Nigeria; human health risk", *Journal of the Nigerian Society of Physical Sciences* **4** (2022) 891. <https://doi.org/10.46481/jnsps.2022.891>.
- [52] G. O. Olutona, "Health risk assessment of heavy metals in sediment of tropical freshwater stream", *Journal of the Nigerian Society of Physical Sciences* **5** (2023) 983. <https://doi.org/10.46481/jnsps.2023.983>.