



Collocation Approach for the Computational Solution Of Fredholm-Volterra Fractional Order of Integro-Differential Equations

G. Ajileye^{a,*}, A.A. James^b, A. M. Ayinde^c, T. Oyedepo^d

^aDepartment of Mathematics and Statistics, Federal University Wukari, Taraba State, Nigeria.

^bDepartment of Mathematics, American University of Nigeria, Yola, Adamawa State, Nigeria.

^cDepartment of Mathematics, Moddibo-Adama University, Yola Nigeria.

^dFederal College of Dental Technology and Therapy, Enugu, Nigeria.

Abstract

In this work, a collocation technique is used to determine the computational solution to fractional order Fredholm-Volterra integro-differential equations with boundary conditions using Caputo sense. We obtained the linear integral form of the problem and transformed it into a system of linear algebraic equations using standard collocation points. The matrix inversion approach is adopted to solve the algebraic equation and substituted it into the approximate solution. We established the uniqueness and convergence of the method and some modelled numerical examples are provided to demonstrate the method's correctness and efficiency. It is observed that the results obtained by our new method are accurate and performed better than the results obtained in the literature. The study will be useful to engineers and scientists. It is advantageous because it addresses the difficulty in tackling fractional order Fredholm-Volterra integro-differential problems using a simple collocation strategy. The approach has the advantage of being more accurate and reducing computer running time.

DOI:10.46481/jnsps.2022.834

Keywords: Collocation, Fredholm-Volterra, Integro-differential, Fractional derivatives, Approximate Solution.

Article History :

Received: 27 May 2022

Received in revised form: 13 August 2022

Accepted for publication: 18 August 2022

Published: 03 October 2022

© 2022 The Author(s). Published by the Nigerian Society of Physical Sciences under the terms of the Creative Commons Attribution 4.0 International license (<https://creativecommons.org/licenses/by/4.0>). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Communicated by: J. Ndam

1. Introduction

Fractional calculus is an aspect in mathematics that studies the properties of integrals combined with noninteger order derivatives. The concept and method of solving differential equations containing fractional derivatives of unknown functions are covered in this field (fractional differential

equations). Many prominent mathematicians, such as Liouville, Grunwald, Riemann, Euler, Langrange, Heaviside, Fourier, Abel, and others, established fractional calculus on a formal foundation[1]. Very recently, scholars developed huge interest in fractional calculus due to its relevance and application to many areas of scientific endeavors. Fractional differential equations, or those containing real and complex order derivatives have been more significant in describing the most broad fields of science and technology with peculiar dynamics of various processes involving complex systems [2].

*Corresponding author tel. no: +2348077092831

Email address: adewa1e.james@aun.edu.ng (G. Ajileye)

Many different approaches have been adopted to investigate the solution of fractional integrodifferential equations, such as Adomian decompositions method [3-5], collocation method [6, 7], Laplace decomposition method [8, 9], Taylor expansion method [10], Least square method [9], differential transform method [11], homotopy perturbation method [12-18], sinc-collocation method [15-17] and variational iteration method [12, 13]. Linna *et al.* [20] considered a numerical method to solve fractional variational problem. They simplified the fractional variational problems by the operational matrices. The operational matrices are based on the Chelyshkov polynomials. The fractional variational problem was transformed into a set of algebraic equations. The unknown coefficients were solved using the Lagrange multiplier techniques. Avcı & Mahmudov [19] proposed a numerical method based on the fractional Taylor vector for solving multi-term fractional differential equations. The main idea of this method was to reduce the given problems to a set of algebraic equations by utilizing the fractional Taylor operational matrix of fractional integration. Some numerical examples were given to demonstrate the accuracy and applicability. The results show that the presented method was efficient and applicable. Fadugba [21] presented the Mellin transform for the solution of the fractional order equations. The Mellin transform approach occurs in many areas of applied mathematics and technology. The Mellin transform of fractional calculus of different flavours; namely the Riemann-Liouville fractional derivative, Riemann-Liouville fractional integral, Caputo fractional derivative and the Miller-Ross sequential fractional derivative were obtained.

In this study, we consider Fredholm-Volterra Integrodifferential equation of fractional order of the the form:

$$\begin{aligned} & D_x^\alpha y(x) - P_1 y'(x) - P_2 y''(x) - P_0 y(x) \\ & = g(x) + \lambda_1 \int_0^x k_1(x,t)y(t) dt \\ & + \lambda_2 \int_0^1 k_2(x,t)y(t) dt, \end{aligned} \quad (1)$$

with the given boundary conditions

$$y(a) = 0, y(b) = 0, a < x < b, \quad (2)$$

where $y(x)$ is to be determined, D_x^α is the Caputo's derivative, $k_1(x,t)$ and

$k_2(x,t)$ are the Fredholm and Volterra integral kernel function respectively. P_j, P_α, λ_j are known constants. $g(x)$ is the known function

2. Basic Definitions

Under this section, we present some definitions and basic concepts of fractional calculus for the formulation of the given problem

Definition 2.1: The Caputo derivative with order $\alpha > 0$ of the given function $f(x)$, $x \in (a, b)$ is defined as [Litfi, Dehghan and Yousefi, 2011]

$${}_x^C D_a^\alpha y(x) = \frac{1}{\Gamma(m-\alpha)} \int_a^x (x-s)^{m-\alpha-1} y^{(m)}(s) ds, \quad (3)$$

where $m-1 \leq \alpha \leq m, m \in \mathbb{N}, x > 0$

Definition 2.2: Let $(a_n), n \geq 0$ be a sequence of real numbers. The power series in x with coefficients a_n is an expression [Edward, Ford and Simpson, 2002]

$$\begin{aligned} y(x) & = a_0 + a_1 x + a_2 x^2 + a_3 x^3 + \dots + a_N x^N \\ & = \sum_{n=0}^N a_n x^n = \phi(x) \mathbf{A}, \end{aligned} \quad (4)$$

where $\phi(x) = [1 \ x \ x^2 \ \dots \ x^N]$, $\mathbf{A} = [a_0 \ a_1 \ \dots \ a_N]^T$, then $y(x, n) = x^n \mathbf{A}$, $n = 0(1)N, n \in \mathbb{Z}^+$.

Definition 2.3: Standard Collocation Method (SCM). This approach is used to find the collocation points that are desired within a certain interval. i.e $[a,b]$ and is given by

$$x_i = a + \frac{(b-a)i}{N}, i = 1, 2, 3, \dots, N. \quad (5)$$

Definition 2.4: A metric on a set M is a function $d : M \times M \rightarrow \mathbb{R}$ with the following properties. For all $x, y \in M$

- (a) $d(x, y) \geq 0$;
- (b) $d(x, y) = 0 \iff x = y$
- (c) $d(x, y) = d(y, x)$
- (d) $d(x, y) \leq d(x, z) + d(x, y)$

If d is a metric on M , then the pair (M, d) is called a metric space.

Definition 2.5: Let (X, d) be a metric space, A mapping $T : X \rightarrow X$ is Lipschitzian if \exists a constant $L > 0$ such that $d(Tx, Ty) \leq Ld(x, y) \forall x, y \in X$.

Definition 2.6: Let $y(x)$ be a continuous function, then

$${}_0 I_x^\beta ({}_0^C D_x^\beta y(x)) = y(x) - \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k, \quad (6)$$

where $m-1 < \beta < 1$.

Let $p(s)$ be an integrable function, then

$${}_0 I_x^\beta (p(s)) = \frac{1}{\Gamma(\beta)} \int_0^x (x-s)^{\beta-1} p(s) ds. \quad (7)$$

3. Mathematical Background

In this section, combination of collocation method and power series approximation is employed for the computational solution of (FVID) of fractional order.

Theorem 3.0: Banach's fixed point theorem Let (X, d) be a complete metric space, then each contraction mapping $T : X \rightarrow X$ has a unique fixed point x of T in X , such that $T(x) = x$

Lemma (3.1). Let $y(x)$ be the solution to (1) subject to (2), the integral form

$$y(x) = W(x) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_2 y''(s)) ds \quad (8)$$

$$\begin{aligned}
& + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_1 y'(s)) ds \\
& + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_0 y(s)) ds \\
& - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left[\lambda_1 \int_0^s k_1(s,t) y(t) dt \right. \\
& \left. + \lambda_2 \int_0^1 k_2(s,t) y(t) dt \right]
\end{aligned}$$

where

$$W(x) = \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k - \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} g(s) ds.$$

Multiply equation (1) by ${}_0I_x^\beta(\cdot)$ gives

$$\begin{aligned}
{}_0I_x^\alpha (D^\alpha y(x)) &= {}_0I_x^\beta (P_2 y''(x)) + {}_0I_x^\beta (P_1 y'(s)) \\
&+ {}_0I_x^\beta (P_0 y(s)) - {}_0I_x^\beta (g(x)) \\
&- {}_0I_x^\beta \left[\lambda_1 \int_0^s k_1(s,t) y(t) dt \right. \\
&\left. + \lambda_2 \int_0^1 k_2(s,t) y(t) dt \right]. \quad (9)
\end{aligned}$$

Using(6) on equation (9)gives

$$\begin{aligned}
y(x) &= \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k + {}_0I_x^\beta (P_2 y''(x)) \\
&+ {}_0I_x^\beta (P_1 y'(s)) + {}_0I_x^\beta (P_0 y(s)) \\
&{}_0I_x^\beta (g(x)) + {}_0I_x^\beta \left(\int_0^b k_1(x,t) y(t) dt \right) \\
&+ {}_0I_x^\beta \left(\int_0^s k_2(s,t) y(t) dt \right)
\end{aligned} \quad (10)$$

Applying equation (4) and (7) to (10) gives

$$\begin{aligned}
y(x) &= \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k \\
&+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} P_2 (\phi(s'')) ds \mathbf{A} \\
&+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} P_1 (\phi(s')) ds \mathbf{A} \\
&+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} P_0 (\phi(s)) ds \mathbf{A} \\
&- \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (g(s)) ds \\
&- \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left[\lambda_1 \int_0^s k_1(s,t) \phi(t) dt \right. \\
&\left. + \lambda_2 \int_0^1 k_2(s,t) \phi(t) dt \right] ds \mathbf{A} \quad (11)
\end{aligned}$$

3.1. Method of Solution

Collocating at x_i in equation (11) gives

$$\begin{aligned}
y(x_i) &= \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k + \\
&\frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_2 (\phi(s'')) ds \mathbf{A} \\
&+ \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_1 (\phi(s')) ds \mathbf{A} \\
&+ \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_0 (\phi(s)) ds \mathbf{A} \\
&- \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} (g(s)) ds - \\
&\frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} \left[\lambda_1 \int_0^s k_1(s,t) \phi(t) dt \right. \\
&\left. + \lambda_2 \int_0^1 k_2(s,t) \phi(t) dt \right] ds \mathbf{A} \quad (12)
\end{aligned}$$

Applying (4) on (12) gives

$$\begin{aligned}
\phi(x_i) \mathbf{A} &= W(x_i) \\
&+ \left[\begin{array}{l} \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_2 (\phi(s'')) ds + \\ \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_1 (\phi(s')) ds + \\ \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_0 (\phi(s)) ds \\ - \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} \left(\lambda_1 \int_0^s k_1(x_i,t) \phi(t) dt \right. \\ \left. + \lambda_2 \int_0^1 k_2(x_i,t) \phi(t) dt \right) ds \end{array} \right] \mathbf{A}, \quad (13)
\end{aligned}$$

where

$$W(x_i) = \sum_{k=0}^N \frac{y^{(k)}(0)}{k!} x^k - \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} g(s) ds.$$

Factorise the values of \mathbf{A} from equation (13) gives

$$W(x_i) = \left[\begin{array}{l} \phi(x_i) - \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_2 (\phi(s'')) ds - \\ \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_1 (\phi(s')) ds - \\ \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_0 (\phi(s)) ds \\ + \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} \\ \left(\lambda_1 \int_0^s k_1(s,t) \phi(t) dt \right. \\ \left. + \lambda_2 \int_0^1 k_2(s,t) \phi(t) dt \right) ds \end{array} \right] \mathbf{A}. \quad (14)$$

Equation 14 can be in the form

$$\tau_n(x_i) \mathbf{A} = W(x_i), \quad (15)$$

where

$$\tau_n(x_i) = \left[\begin{array}{l} \phi(x_i) - \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_2 (\phi(s'')) ds - \\ \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_1 (\phi(s')) ds - \\ \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} P_0 (\phi(s)) ds + \frac{1}{\Gamma(\alpha)} \int_0^{x_i} (x_i-s)^{\alpha-1} \\ \left(\lambda_1 \int_0^s k_1(s,t) \phi(t) dt + \lambda_2 \int_0^1 k_2(s,t) \phi(t) dt \right) ds, \end{array} \right]$$

and

$$\mathbf{A} = [a_0 \ a_1 \ \dots \ a_N]^T.$$

Multiply both sides of equation(15) by $\tau_n^{-1}(x_i)$ gives

$$\mathbf{A} = \tau_n^{-1}(x_i) W(x_i). \quad (16)$$

Substituting \mathbf{A} into the approximate solution (4) gives

$$y(x) = \phi(x_i) \tau_n^{-1}(x_i) W(x_i) \quad (17)$$

4. Uniqueness of the Method

In order to establish the uniqueness of the method, we introduce the following hypothesis

$$H_1 : k_1^* = \max_{x \in [0,1]} \int_0^x \lambda_1 |k_1(x, t)| dt$$

$$H_2 : k_2^* = \max_{x \in [0,1]} \int_0^1 \lambda_2 |k_2(x, t)| dt$$

$$H_3 : |y_1^{(m)} - y_2^{(m)}| \leq L_m |y_1 - y_2|.$$

Lemma (4.1) (q-contraction) Let $T : X \rightarrow X$ be a mapping defined by theorem (3.0) for $y_1, y_2 \in X$. T is q -contraction if and only if

$$\left[\frac{\sum_{n=0}^2 L_n - (k_1^* + k_2^*)}{\Gamma(\alpha + 1)} \right] < 1,$$

then there exist a unique solution of T .

$$\begin{aligned} (Ty_1)(x) &= W(x) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_2 y_1''(s)) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_1 y_1'(s)) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_0 y_1(s)) ds \\ &- \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left[\lambda_1 \int_0^s k_1(s, t) y_1(t) dt \right. \\ &\left. + \lambda_2 \int_0^1 k_2(s, t) y_1(t) dt \right] ds \end{aligned}$$

and

$$\begin{aligned} (Ty_2)(x) &= W(x) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_2 y_2''(s)) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_1 y_2'(s)) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_0 y_2(s)) ds \\ &- \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left[\lambda_1 \int_0^s k_1(s, t) y_2(t) dt \right. \\ &\left. + \lambda_2 \int_0^1 k_2(s, t) y_2(t) dt \right] ds. \end{aligned}$$

Thus,

$$\begin{aligned} &|(Ty_1)(x) - (Ty_2)(x)| \\ &= \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} P_2 |y_1''(s) - y_2''(s)| ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} P_1 |y_1'(s) - y_2'(s)| ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} P_0 |y_1(s) - y_2(s)| ds \\ &- \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left[\int_0^s \lambda_1 |k_1(s, t)| |y_1(t) - y_2(t)| dt \right. \\ &\left. + \int_0^1 \lambda_2 |k_2(s, t)| |y_1(t) - y_2(t)| dt \right] ds. \end{aligned}$$

Taking maximum of both sides and using H_1 to H_3 gives

$$d(Ty_1(x), Ty_2(x)) \leq \left[\frac{L_2 + L_1 + L_0 - (k_1^* + k_2^*)}{\Gamma(\alpha + 1)} \right] d(y_1, y_2)$$

Since T is a contraction

$$\left[\frac{\sum_{n=0}^2 L_n - (k_1^* + k_2^*)}{\Gamma(\alpha + 1)} \right] < 1.$$

We can conclude that T has a unique solution.

5. Convergence Analysis

We establish the convergence of the method

$$\begin{aligned} y_N(x) &= W(x) + \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_2 y_N''(s)) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_1 y_N'(s)) ds \\ &+ \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} (P_0 y_N(s)) ds \\ &- \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \left[\lambda_1 \int_0^s k_1(s, t) y_N(t) dt \right. \\ &\left. + \lambda_2 \int_0^1 k_2(s, t) y_N(t) dt \right] ds. \end{aligned} \quad (18)$$

Subtract (8) from (18) gives

$$E_N(x) = y_N(x) - y(x).$$

Hence

$$\begin{aligned} |E_N(x)| &= \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} |E_N(s) (P_2 + P_1)| ds \\ &- \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \\ &\left[|\lambda_1| \left| \int_0^s k_1(s, t) E_N(t) dt \right| \right. \\ &\left. + |\lambda_2| \left| \int_0^1 k_2(s, t) E_N(t) dt \right| \right] ds. \end{aligned}$$

(20)

Therefore

$$\begin{aligned} \frac{\|E_N(x)\|_\infty}{\|E_N(t)\|_\infty} &\leq \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \\ &\times \left[\begin{aligned} &|E_n(s) (P_2 + P_1)| \\ &- \frac{1}{\Gamma(\alpha)} \int_0^x (x-s)^{\alpha-1} \\ &\left(|\lambda_1| \left| \int_0^s k_1(s, t) E_n(t) dt \right| \right. \\ &\left. + |\lambda_2| \left| \int_0^1 k_2(s, t) E_n(t) dt \right| \right) \end{aligned} \right] ds. \end{aligned}$$

The method converges

6. Numerical Examples

In this section, two numerical problems with boundary conditions are presented to test the efficiency and simplicity of the method. All computations are done with the aid of Maple 18. Let $y_n(x)$ and $y(x)$ be the approximate and exact solution respectively. $Error_N = |y_n(x) - y(x)|$

Example 1:

Considering fractional integro-differential equation [17]

$$y''(x) + \frac{1}{x} D_x^{0.5} y(x) + \frac{1}{x^2} y(x) = f(x) + \int_0^x \sin(x-t)y(t)dt + \int_0^1 \cos(x-t)y(t)dt,$$

subject to the boundary conditions

$$y(0) = 0, y(1) = 0,$$

where

$$f(x) = 5 + 1.50451x^{0.5} - 13x - 1.80541x^{1.5} - x^2 + x^3 - 2.067x \cos(x) + 5.95385 \sin(x).$$

Exact solution $y(x) = x^2(1-x)$

Solution 1

We solve this problem at $N = 3$ and 5 but we use $N = 3$ for demonstration. Integral form of example 1 is

$$y''(x) + \frac{1}{x} \left(\frac{1}{\Gamma(1-0.5)} \int_0^x (x-t)^{-0.5} y^{(1)}(t)dt \right) + \frac{1}{x^2} y(x) = f(x) + \int_0^x \sin(x-t)y(t)dt + \int_0^1 \cos(x-t)y(t)dt \quad (21)$$

Using approximate solution (4) on (14) gives

$$\begin{bmatrix} \phi''(x) + \frac{1}{x} \left(\frac{1}{\Gamma(1-0.5)} \int_0^x (x-t)^{-0.5} \frac{d}{dx}(\phi(t))dt \right) + \frac{1}{x^2} \phi''(x) - \int_0^x \sin(x-t)\phi(t)dt - \int_0^1 \cos(x-t)\phi(t)dt \\ \mathbf{A} = f(x) \end{bmatrix} \quad (22)$$

Equation (15) gives

$$\tau(x)\mathbf{A} = f(x), \quad (23)$$

where

$$\tau(x) = \phi''(x) + \frac{1}{x} \left(\frac{1}{\Gamma(1-0.5)} \int_0^x (x-t)^{-0.5} \frac{d}{dx}(\phi(t))dt \right) + \frac{1}{x^2} \phi''(x) - \int_0^x \sin(x-t)\phi(t)dt - \int_0^1 \cos(x-t)\phi(t)dt.$$

Collocating at $x_3 = \left[\frac{1}{3} \quad \frac{2}{3} \quad 1 \right]$ and substituting the boundary conditions gives

$$\tau(x)^* \mathbf{A} = f(x)^*, \quad (24)$$

where

$$\tau_i(x) = \begin{bmatrix} 8.1837497830 & 4.5298367530 & 3.5647166670 & 2.4321976680 \\ 1.0664602830 & 2.3357941530 & 3.8888778550 & 5.4288519160 \\ -0.3011686789 & 1.5101524580 & 4.1068429140 & 8.5147669310 \\ 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix}$$

Table 1. Exact and approximate values, Example 1

x	Exact	N = 3	N = 5
0.2	0.032000000000	0.031999821270	0.031999851830
0.4	0.096000000000	0.095999891010	0.095999961860
0.6	0.144000000000	0.143999795600	0.143999892900
0.8	0.128000000000	0.127999546700	0.127999648400
1.0	0.000000000000	-8.444000000000e-7	-7.382000000000e-7

Table 2. Absolute Error for Example 1

x	error ₃	error ₅	error _{[17]=32}
0.2	-1.787300000000e-7	-1.481700000000e-7	2.048e-5
0.4	-1.089900000000e-7	-3.814000000000e-8	2.503e-5
0.6	-2.044000000000e-7	-1.071000000000e-8	1.789e-5
0.8	-1.071000000000e-7	-3.516000000000e-8	7.682e-6
1	-8.444000000000e-7	-7.382000000000e-8	3.034e-6

$$f(x) = [1.1325152410 \quad -1.5399784720 \quad -4.4079289640 \quad 0 \quad 0] .$$

We now solve for the unknown values \mathbf{A} (17) making use of matrix inversion results in;

$$y_3 = \begin{pmatrix} -4.2497275388 \times 10^{-7} + 0.16917e - 5x + \\ 0.9999976497x^2 - 0.999997608x^3 \end{pmatrix} .$$

Applying the same procedure for $N = 5$ gives

$$y_5 = \begin{pmatrix} -3.4773620430 \times 10^{-7} + 8.9350760390 \times 10^7 x + 1.0000017198x^2 \\ -1.0000070682x^3 + 0.57092e - 5x^4 - 0.16454e - 5x^5 \end{pmatrix}$$

Example 2

Considering fractional integro-differential equation [17]

$$y''(x) + D_x^\alpha y(x) = f(x) + 2 \int_0^x (x-t)y(t)dt + \int_0^1 (x^2-t)y(t)dt$$

with the given boundary conditions

$$y(0) = 0, y(1) = 0$$

where

$$f(x) = -\frac{1}{30} - 6x + \frac{181x^2}{20} + 4x^3 - \frac{x^5}{10} + \frac{x^6}{15} \text{ and } \alpha = 1$$

Exact solution $y(x) = x^3(x-1)$

Solution 2

We solve this problem at $N = 4$ and 5 , however, make use of $N = 4$ for demonstration.

Integral form of example 2 is

$$y''(x) + \left(\frac{1}{\Gamma(2-1)} \int_0^x (x-t)^0 y^{(2)}(t)dt \right) = f(x) + 2 \int_0^x (x-t)y(t)dt + \int_0^1 (x^2-t)y(t)dt \quad (25)$$

Table 3. Exact and approximate values, Example 2

x	Exact	$N = 4$	$N = 5$
0.2	-0.6400000000e-2	-0.6400000000e-2	-0.6400000000e-2
0.4	-0.3840000000e-1	-0.3840000000e-1	-0.3840000000e-1
0.6	-0.8640000000e-1	-0.8640000000e-1	-0.8640000000e-1
0.8	-0.1024000000	-0.1024000000	-0.1024000000
1.0	-0.9000000000e-3	-0.9000000000e-3	-0.9000000000e-3

Table 4. Absolute Error for Example 2

x	error ₄	error ₅	error _{[17]=64}
0.2	0.0	0.0	2.931e-7
0.4	0.0	0.0	3.915e-7
0.6	0.0	0.0	2.696e-7
0.8	0.0	0.0	1.011e-7
1	0.0	0.0	3.596e-8

Using approximate solution (4) on (18) gives

$$\begin{bmatrix} \phi''(x) + \left(\frac{1}{\Gamma(2-1)} \int_0^x (x-t)^0 \frac{d^2}{dx^2}(\phi(t))dt \right) \\ -2 \int_0^x (x-t) \phi(t)dt - \int_0^1 (x^2-t) \phi(t) dt \end{bmatrix} \mathbf{A} = f(x). \quad (26)$$

Equation (6.6) gives

$$\tau(x)\mathbf{A} = f(x) \quad (27)$$

where

$$\tau(x) = \begin{bmatrix} \phi''(x) + \left(\frac{1}{\Gamma(2-1)} \int_0^x (x-t)^0 \frac{d^2}{dx^2}(\phi(t))dt \right) \\ -2 \int_0^x (x-t) \phi(t)dt - \int_0^1 (x^2-t) \phi(t) dt \end{bmatrix}$$

Collocating at $x_4 = \left[0 \quad \frac{1}{4} \quad \frac{2}{4} \quad \frac{3}{4} \quad 1 \right]$ and substituting the boundary conditions gives

$$\tau(x)^* \mathbf{A} = f(x)^* \quad (28)$$

where

$$\tau_i(x) = \begin{bmatrix} \frac{1}{2} & 0.3333333333 & 2.250000000 & 0.200000000 & 0.1666666667 \\ \frac{1}{2} & \frac{59}{192} & \frac{4193}{305} & \frac{19169}{10240} & \frac{59393}{61440} \\ \frac{1}{2} & \frac{1}{4} & \frac{96}{1851} & \frac{64}{64211} & \frac{960}{522457} \\ \frac{1}{2} & \frac{37}{192} & \frac{517}{29} & \frac{10240}{181} & \frac{522457}{61440} \\ \frac{1}{2} & \frac{1}{6} & \frac{12}{0} & \frac{20}{0} & \frac{30}{0} \\ \frac{1}{2} & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 \end{bmatrix}$$

$$f(x) = \begin{bmatrix} -\frac{1}{30} & -\frac{55621}{61440} & -\frac{131}{480} & \frac{137191}{61440} & \frac{419}{60} & 0 & 0 \end{bmatrix}$$

We solve for the unknown values \mathbf{A} using matrix inversion, results;

$$y_4 = \begin{pmatrix} -2.2211399390 \times 10^{-13} \\ +3.0783153800 \times 10^{-12}x \\ -1.4125589590 \times 10^{-11}x^2 \\ -1.0000000000x^3 + 1.0000000000x^4 \end{pmatrix}$$

Applying the same procedure for $N = 5$ gives

$$y_5 = \begin{pmatrix} 5.329070518000 \times 10^{-15} \\ +1.781685910000 \times 10^{-12}x \\ -1.580957587000 \times 10^{-11}x^2 \\ -1.000000000000x^3 \\ +1.000000000000x^4 \\ +1.062971933000 \times 10^{-11}x^5 \end{pmatrix}$$

7. Conclusion

Collocation approach is utilized to solving the fractional order Fredholm-Volterra integro-differential problems in this paper. The applied method is consistent, efficient and easy to compute. The results of the numerical example 1 as shown in table 1 shows that the approximate solution at $N = 3$ gives $y_3(x) = -4.2497275388 \times 10^{-7} + 0.16917e - 5x + 0.9999976497x^2 - 0.999997608x^3$ and solving at $N = 5$ indicates that as the value of N increases, the error becomes smaller. We also compare our absolute errors with Alkan *et al* (2017) as shown in table 2, this clearly shows that our method performs better. The results of the numerical example 2 as shown in table 3 shows that the approximate solution obtained at $N = 4$ and 5 converges to the exact solution. Table 4 compares the errors in the method proposed by Alkan *et al* (2017) at $N = 64$ and errors in our new method which converges to the exact solution at $N = 4$.

References

- [1] P. Kamal, *Higher order numerical methods for fractional order differential equations*, Doctoral dissertation, University of Chester, United Kingdom, 2015. <http://hdl.handle.net/10034/613354>.
- [2] D. Baleanu, K. Diethelm & J. J. Trujillo, *Fractional calculus: Models and numerical methods*, Series on complexity nonlinearity and chaos, World Scientific 2012.
- [3] R. H. Khan & H. O. Bakodah, "Adomian decomposition method and its modification for nonlinear Abel's integral equations", *Computers and Mathematics with Applications* 7 (2013) 2349.
- [4] R. C. Mittal & R. Nigam, "Solution of fractional integro-differential equations by Adomian decomposition method", *The International Journal of Applied Mathematics and Mechanics* 2 (2008) 87.
- [5] S. Momani & M. A. Noor, "Numerical methods for fourth-order fractional integro-differential equations", *Applied Mathematics and Computation* 1 (2006) 754.
- [6] A. O. Agbolade & T. A. Anake, "Solution of first order volterra linear integro differential equations by collocation method", *J. Appl. Math.*, Article ID. 1510267 (2017). doi:10.1155/2017/1510267
- [7] O. Cem, S. Mehmet & D. O. Arzu, "Chelyshkov collocation approach to solving the systems of linear functional differential equations", *New Trend in Mathematical Sciences (NTMSCI)* 4 (2015) 83.
- [8] D. Nazari & M. Jahanshahi, "Some notes on multi-order fractional integro-differential equations", *43rd Annual Iranian Mathematics Conference* (2012) pp 27.
- [9] D. S. Mohammed, "Numerical solution of fractional integro-differential equations by least square method and shifted Chebyshev polynomial", *Mathematical Problems in Engineering* 2014 (2014) 431965. <https://doi.org/10.1155/2014/431965>
- [10] L. Huang, X. F. Li, Y. Zhao & X. Y. Duan, "Approximate solution of fractional integro-differential equations by Taylor expansion method", *Computers & Mathematics with Applications* 3 (2011) 1127.
- [11] A. Arikoglu & I. Ozkol, "Solution of fractional integro-differential equations by using fractional differential transform method", *Chaos, Solitons & Fractals* 2 (2009) 521.
- [12] A. A. Elbeleze, A. Klman & B. M. Taib, "Approximate solution of integro-differential equation of fractional (arbitrary) order", *Journal of King Saud University - Science* 28 (2016) 61.
- [13] Y. Nawaz, "Variational iteration method and homotopy perturbation method for fourth-order fractional integro-differential equations", *Computers & Mathematics with Applications* 8 (2011) 2330.
- [14] R. K. Saeed & H. M. Sdeq, "Solving a system of linear fredholm fractional integro-differential equations using homotopy perturbation method", *Australian Journal of Basic and Applied Sciences* 4 (2010) 633.
- [15] S. Alkan, "A new solution method for nonlinear fractional integro-differential equations", *Discrete and Continuous Dynamical Systems - Series S* 6 (2015) 1065.

- [16] S. Alkan & A. Secer, "Solution of nonlinear fractional boundary value problems with nonhomogeneous boundary conditions", *Applied and Computational Mathematics* **3** (2015).
- [17] S. Alkan & V. F. Hatipoğlu, "Approximate solutions of Volterra-Fredholm integro-differential equations of fractional order", *Tbilisi Mathematical Journal* **2** (2017) 1.
- [18] O. A. Uwaheren, A. F. Adebisi & O. A. Taiwo, "Perturbed Collocation Method For Solving Singular Multi-order Fractional Differential Equations of Lane-Emden Type", *Journal of the Nigerian Society of Physical Sciences* **3** (2020) 141. <https://doi.org/10.46481/jnsps.2020.69>.
- [19] I. Avcı & N. I. Mahmudov, "Numerical Solutions for Multi-Term Fractional Order Differential Equations with Fractional Taylor Operational Matrix of Fractional Integration", *Mathematics* **1** (2020) 96. <https://doi.org/10.3390/math8010096>.
- [20] L. Linna, W. Zhirou & H. Qiongdan, "A numerical method for solving feactional variational problems by operational matrix based on Chelyshekov polynomial", *Engineering Letters* **28** (2020) 486.
- [21] S. E. Fadugba, "Solution of Fractional Order Equations in the Domain of the Mellin Transform", *Journal of the Nigerian Society of Physical Sciences* **4** (2019) 138. <https://doi.org/10.46481/jnsps.2019.31>