



# Masses and thermal properties of a Charmonium and Bottomonium Mesons

E. P. Inyang<sup>a,\*</sup>, E. O. Obisung<sup>c</sup>, P. C. Iwuji<sup>c</sup>, J. E. Ntibi<sup>b</sup>, J. Amajama<sup>c</sup>, E. S. William<sup>b</sup>

<sup>a</sup> Department of Physics, National Open University of Nigeria, Jabi, Abuja, Nigeria

<sup>b</sup> Theoretical Physics Group, Department of Physics, University of Calabar, PMB 1115, Calabar, Nigeria

<sup>c</sup> Department of Physics, University of Calabar, PMB 1115, Calabar, Nigeria

## Abstract

In this research, we model Hulthén plus generalized inverse quadratic Yukawa potential to interact in a quark-antiquark system. The solutions of the Schrödinger equation are obtained using the Nikiforov-Uvarov method. The energy spectrum and normalized wave function were obtained. The masses of the heavy mesons for different quantum states such as 1S, 2S, 1P, 2P, 3S, 4S, 1D, and 2D were predicted as 3.096 GeV, 3.686 GeV, 3.327 GeV, 3.774 GeV, 4.040 GeV, 4.364 GeV, 3.761 GeV, and 4.058 GeV respectively for charmonium ( $c\bar{c}$ ). Also, for bottomonium ( $b\bar{b}$ ) we obtained 9.460 GeV, 10.023 GeV, 9.841 GeV, 10.160 GeV, 10.345 GeV, 10.522 GeV, and 10.142 GeV for different states of 1S, 2S, 1P, 2P, 3S, 4S, 1D respectively. The partition function was calculated from the energy spectrum, thereafter other thermal properties were obtained. The results obtained showed an improvement when compared with the work of other researchers and excellently agreed with experimental data with a percentage error of 1.60 % and 0.46 % for ( $c\bar{c}$ ) and ( $b\bar{b}$ ), respectively.

DOI:10.46481/jnsps.2022.884

**Keywords:** Schrödinger equation; Nikiforov-Uvarov method; heavy mesons; Thermal properties; Hulthén plus generalized inverse quadratic Yukawa potential.

## Article History :

Received: 23 June 2022

Received in revised form: 9 July 2022

Accepted for publication: 11 July 2022

Published: 19 August 2022

© 2022 The Author(s). Published by the Nigerian Society of Physical Sciences under the terms of the Creative Commons Attribution 4.0 International license (<https://creativecommons.org/licenses/by/4.0>). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Communicated by: C. Edet, P. Amadi & R. Khordad

## 1. Introduction

In non-relativistic quantum mechanics, researchers work mostly on the solutions of the Schrödinger equation (SE) to obtain the eigenvalues and eigenfunction. These eigenvalues and eigenfunction can be used to study the mass spectra (MS) of the heavy mesons, and theoretic measures, among other things [1, 2, 3, 4]. In describing the interaction of charmonium

( $c\bar{c}$ ) and bottomonium ( $b\bar{b}$ ), a Cornell potential (CP) is used because of the Coulomb interaction and a confining term [5]. More so, in solving the SE with any potential of choice, analytical methods are employed. Most of the frequently used analytical methods are as follows, the Nikiforov-Uvarov (NU) method [6-12], the Nikiforov-Uvarov Functional Analysis (NUFA) method [13-16], the series expansion method (SEM) [17, 18, 19], Laplace transformation method (LTM) [20], Asymptotic Iteration Method [21], WKB approximation method [22-24], among others [25-31]. The investigation of the MS with CP has gained remarkable interest, and has

\*Corresponding author tel. no: +2348032738073

Email address: [etidophysics@gmail.com](mailto:etidophysics@gmail.com), [einyang@noun.edu.ng](mailto:einyang@noun.edu.ng)  
(E. P. Inyang )

attracted the attention of many scholars [32-35]. For instance, Mutuk [36] solved the SE with CP using a neural network approach. The bottomonium, charmonium, and bottom-charmed spin-averaged spectra were calculated. Using, the vibrational method (VM) and supersymmetric quantum mechanics Vega and Flores, [37] obtained the analytical solutions of the SE with CP. The eigenvalues were used to calculate the MS of the HMs. Also, Kumar et al., [38] used the NUFA method to solve the SE with generalized Cornell potential. The result was used to predict the MS of the HMs. Furthermore, Hassanabadi et al. [39], used the VM to solve the SE with CP. The eigenvalues were used to calculate the mesonic wave function. In recent times, the study of MS of the HMs with exponential-type potentials has aroused the interest of scholars [40, 41]. Potential models such as Yukawa potential [42], Varshni [43], screened Kratzer potential [44], and so on have been used in the prediction of the MS of the HMs. For instance, Purohit et al [45] combined linear plus modified Yukawa potential to obtain the masses of the HMs through the solutions of the Klein-Gordon equation. The SE for most of the potentials with spin addition cannot be solved analytically; hence, numerical solutions such as Runge-Kutte approximation [46], Numerov matrix method [47], Fourier grid Hamiltonian method [48], and so on [49] are employed. Also, adding spin enables one to determine other properties of the mesons like decay properties and root mean square radii. However, we have considered our mesons as spinless particles for easiness [1, 40, 50, 51, 52]. Recently, researchers have studied thermal properties (TPs) of the HMs [40, 53]. For instance, Abu-Shady et al., [54] studied the TPs of HMs with extended Cornell potential. Also Inyang et al., [55] obtained the TPs of the HMs with exponential potential. Abu-Shady and Ezz-Alarab [40], solved the N-radial SE with the Trigonometric Rosen-Morse potential, and the obtained energy equation was used in studying TPs and masses of heavy and heavy-light mesons.

Cao et al., [56], studied the TPs of light mesons, including the temperature dependence of their masses by solving the spatial correlation. In recent times scholars, have proposed new potentials through the combination of two potentials [57, 58]. It's observed that the combination of at least two potential models has a propensity to fit experimental data more than a single potential [55]. Okoi et al.,[59] proposed the Hulthén plus generalized inverse quadratic Yukawa potential (HGIQYP) to obtain the bound state energy and expectation values. Hulthén potential [60] and generalized inverse quadratic Yukawa potential [61], is applied to diverse areas of physics such as nuclear and particle and so on.

The aim of this study is in three folds. First, the HGIQYP will be modeled to fit in the CP, secondly, the SE will be solved using the well-known NU method, and finally to predict the masses of charmonium and bottomonium mesons and calculate its TPs.

The HGIQYP takes the form [60]:

$$V(q) = -\frac{R_0 e^{\theta q}}{1 - e^{\theta q}} - R_1 \left(1 + \frac{e^{\theta q}}{q}\right)^2 \quad (1)$$

where  $R_0$  and  $R_1$  are the strength of the potential,  $\theta$  is the screening parameter and  $q$  is inter-nuclear distance. We model equation (1) to interact in the quark-antiquark system by expanding the exponential terms in a series of powers up to order three and have,

$$V(q) = -\frac{G_0}{q} + G_1 q + G_2 q^2 + \frac{G_3}{q^2} + G_4 \quad (2)$$

where

$$G_0 = \frac{R_0}{\sigma} + 2R_1 - 2R_1\alpha, \quad G_1 = -\frac{\alpha R_0}{6} - R_1\alpha^2 \quad (3)$$

$$G_2 = \frac{R_1\alpha^3}{3}, \quad G_3 = -R_1, \quad G_4 = \frac{R_0}{2} - R_1 + 2\alpha R_1 - 2R_1\alpha^2$$

## 2. The solutions of the Schrödinger equation with HGIQYP

In this research, the NU method is employed. The details are found in [62]. The SE is given as [63]

$$\frac{d^2 U(q)}{dq^2} + \left[ \frac{2\mu}{\hbar^2} (E_{nl} - V(q)) - \frac{\ell(\ell + 1)}{q^2} \right] U(q) = 0 \quad (4)$$

where  $\ell$  is the angular momentum quantum number,  $\mu$  is the reduced mass for the quark-antiquark particle,  $q$  is the inter-particle distance, and  $\hbar$  is reduced plank constant. Then, we substitute equation (2) into equation (4), and get

$$\frac{d^2 U(q)}{dq^2} + \left[ \frac{2\mu E_{nl}}{\hbar^2} + \frac{2\mu G_0}{\hbar^2 q} - \frac{2\mu G_1 q}{\hbar^2} - \frac{2\mu G_2 q^2}{\hbar^2} - \frac{2\mu G_3}{\hbar^2 q^2} - \frac{2\mu G_4}{\hbar^2} - \frac{\ell(\ell + 1)}{q^2} \right] U(q) = 0 \quad (5)$$

Transformation of  $q$  (in equation (8) to  $w$  coordinates yields equation (6),

$$w = \frac{1}{q}, \quad q > 0 \quad (6)$$

The second derivative of equation (6) is given as

$$\frac{d^2 U(q)}{dq^2} = 2w^3 \frac{dU(w)}{dw} + w^4 \frac{d^2 U(w)}{dw^2} \quad (7)$$

Replacing equations (6) and (7) in equation (8) gives;

$$\frac{d^2 U(w)}{dw^2} + \frac{2w}{w^2} \frac{dU}{dw} + \frac{1}{w^4} \left[ \frac{2\mu E_{nl}}{\hbar^2} + \frac{2\mu G_0 w}{\hbar^2} - \frac{2\mu G_1}{\hbar^2 w} - \frac{2\mu G_2}{\hbar^2 w^2} - \frac{2\mu G_3 w^2}{\hbar^2} - \frac{2\mu G_4}{\hbar^2} - l(l + 1) w^2 \right] U(w) = 0 \quad (8)$$

The approximation scheme (AS) is introduced by assuming that there is a characteristic radius  $r_0$  of the meson. The AS is achieved by the expansion of  $\frac{G_1}{w}$  and  $\frac{G_1^2}{w^2}$  in a power series

around  $r_0$  ; i.e. around  $\phi \equiv \frac{1}{r_0}$ , up to the second-order [64].

By setting  $y = w - \phi$  and around  $y = 0$  we expand it in powers of series as:

$$\frac{G_1}{w} = \frac{G_1}{y + \phi} = \frac{G_1}{\phi \left(1 + \frac{y}{\phi}\right)} = \frac{G_1}{\phi} \left(1 + \frac{y}{\phi}\right)^{-1} \tag{9}$$

Equation (9) yields

$$\frac{G_1}{w} = G_1 \left( \frac{3}{\phi} - \frac{3z}{\phi^2} + \frac{z^2}{\phi^3} \right) \tag{10}$$

Similarly,

$$\frac{G_2}{w^2} = G_2 \left( \frac{6}{\phi^2} - \frac{8w}{\phi^3} + \frac{3w^2}{\phi^4} \right) \tag{11}$$

Placing equations (10) and (11) into equation (13), we obtain:

$$\frac{d^2 U(w)}{dw^2} + \frac{2w}{w^2} \frac{dU(w)}{dw} + \frac{1}{w^4} \left[ -\varepsilon + X_0 w - X_1 w^2 \right] U(w) = 0 \tag{12}$$

where

$$-\varepsilon = \left( \frac{2\mu E_{nl}}{\hbar^2} - \frac{6\mu G_1}{\hbar^2 \phi} - \frac{12\mu G_2}{\hbar^2 \phi^2} - \frac{2\mu G_4}{\hbar^2} \right), \tag{13}$$

$$X_0 = \left( \frac{2\mu G_0}{\hbar^2} + \frac{6\mu G_1}{\hbar^2 \phi^2} + \frac{16\mu G_2}{\hbar^2 \phi^3} \right)$$

$$X_1 = \left( \frac{2\mu G_1}{\hbar^2 \phi^3} + \frac{6\mu G_2}{\hbar^2 \phi^4} + \frac{2\mu G_3}{\hbar^2} + \gamma \right), \gamma = l(l+1)$$

Connecting equation (12) and equation (1) of [62], we obtain

$$\tilde{\tau}(w) = 2w, \sigma(w) = z^2 \tag{14}$$

$$\tilde{\sigma}(w) = -\varepsilon + \alpha w - \beta w^2$$

$$\sigma'(w) = 2w, \sigma''(w) = 2$$

Plugging equation (23) into equation (8) of [56], gives

$$\pi(w) = \pm \sqrt{\varepsilon - X_0 w + (X_1 + k) w^2} \tag{15}$$

To determine  $k$ , in equation (15), the discriminant of the function equation (16) and equation (17) are obtained as

$$k = \frac{X_0^2 - 4X_1 \varepsilon}{4\varepsilon} \tag{16}$$

$$\pi(w) = \pm \left( \frac{X_0 w}{2\sqrt{\varepsilon}} - \frac{\varepsilon}{\sqrt{\varepsilon}} \right) \tag{17}$$

For a physically acceptable solution, the negative part of equation (17) is required, upon differentiating we get.

$$\pi'_-(w) = -\frac{X_0}{2\sqrt{\varepsilon}} \tag{18}$$

Placing equation (23) and equation (17) into equation (6) of Ref. [62] gives

$$\tau(w) = 2w - \frac{X_0 w}{\sqrt{\varepsilon}} + \frac{2\varepsilon}{\sqrt{\varepsilon}} \tag{19}$$

Equation (19) gives

$$\tau'(w) = 2 - \frac{X_0}{\sqrt{\varepsilon}} \tag{20}$$

Using equations (19) and equation (21) of Ref. [62], we have the following,

$$\lambda = \frac{X_0^2 - 4X_1 \varepsilon}{4\varepsilon} - \frac{X_0}{2\sqrt{\varepsilon}} \tag{21}$$

$$\lambda_n = \frac{nX_0}{\sqrt{\varepsilon}} - n^2 - n \tag{22}$$

Equating equations (21) and (21), followed by the substitution of equations (5) and (14) yielded the energy spectrum of the HGIQYP

$$E_{nl} = \frac{R_0}{2} - R_1 + 2\vartheta R_1 - 2R_1 \vartheta^2 - \frac{\vartheta R_0}{2\phi} - \frac{3R_1 \vartheta^2}{\phi} + \frac{2R_1 \vartheta^3}{\phi^2} \tag{23}$$

$$- \frac{\hbar^2}{8\mu} \left[ \frac{\frac{6\mu}{\hbar^2 \phi^2} \left( -\frac{\vartheta R_0}{6} - R_1 \vartheta^2 \right) + \frac{2\mu}{\hbar^2} \left( \frac{R_0}{\vartheta} + 2R_1 - 2R_1 \vartheta \right) + \frac{16\mu R_1 \vartheta^3}{3\hbar^2 \phi^3}}{n + \frac{1}{2} + \sqrt{\left( \frac{1}{2} + l \right)^2 + \frac{2\mu}{\hbar^2 \phi^3} \left( -\frac{\vartheta R_0}{6} - R_1 \vartheta^2 \right) + \frac{2\mu R_1 \vartheta^3}{\hbar^2 \phi^4} - \frac{2\mu R_1}{\hbar^2}} \right]^2$$

The wave function, is obtained by putting equations (23) and (17) into equation (4) of Ref. [62]

$$\frac{d\phi}{\phi} = \left( \frac{\varepsilon}{w^2 \sqrt{\varepsilon}} - \frac{X_0}{2w \sqrt{\varepsilon}} \right) dw \tag{24}$$

Integration of equation (24) gives

$$\phi(w) = w^{-\frac{X_0}{2\sqrt{\varepsilon}}} e^{-\frac{\varepsilon}{w\sqrt{\varepsilon}}} \tag{25}$$

Putting equations (23) and (19) into equation (26) of Ref. [62] we have

$$\rho(w) = w^{-\frac{X_0}{\sqrt{\varepsilon}}} e^{-\frac{2\varepsilon}{w\sqrt{\varepsilon}}} \tag{26}$$

The substitution of equations (23) and (26) into equation (5) of Ref.[62] gives

$$y_n(w) = B_n e^{\frac{2\varepsilon}{w\sqrt{\varepsilon}}} z^{\frac{X_0}{\sqrt{\varepsilon}}} \frac{d^n}{dw^n} \left[ e^{-\frac{2\varepsilon}{w\sqrt{\varepsilon}}} w^{n-\frac{X_0}{\sqrt{\varepsilon}}} \right] \tag{27}$$

The Rodrigues' formula of the associated Laguerre polynomials is

$$L_n^{\frac{X_0}{\sqrt{\varepsilon}}} \left( \frac{2\varepsilon}{w\sqrt{\varepsilon}} \right) = \frac{1}{n!} e^{\frac{2\varepsilon}{w\sqrt{\varepsilon}}} w^{\frac{X_0}{\sqrt{\varepsilon}}} \frac{d^n}{dw^n} \left( e^{-\frac{2\varepsilon}{w\sqrt{\varepsilon}}} w^{n-\frac{X_0}{\sqrt{\varepsilon}}} \right) \tag{28}$$

where  $B_n = \frac{1}{n!}$ . Hence,

$$y_n(w) \equiv L_n^{\frac{X_0}{\sqrt{\varepsilon}}} \left( \frac{2\varepsilon}{w\sqrt{\varepsilon}} \right) \quad (29)$$

The substitution of equations (25) and (29) into equation (2) of Ref. [62], gives the wave function in terms of associated Laguerre polynomials as

$$\psi(w) = N_{nl} w^{-\frac{X_0}{2\sqrt{\varepsilon}}} e^{-\frac{\varepsilon}{w\sqrt{\varepsilon}}} L_n^{\frac{X_0}{\sqrt{\varepsilon}}} \left( \frac{2\varepsilon}{w\sqrt{\varepsilon}} \right) \quad (30)$$

where  $N_{nl}$  is normalization constant, which can be obtained from

$$\int_0^\infty |\psi_{nl}(r)|^2 dr = 1 \quad (31)$$

Inserting equation (30) into (31) with  $w = 1/r$  gives

$$N_{nl}^2 \int_0^\infty r^{X_0/\sqrt{\varepsilon}} e^{-2\sqrt{\varepsilon}r} \left[ L_n^{X_0/\sqrt{\varepsilon}} (2\sqrt{\varepsilon}r) \right]^2 dr = 1 \quad (32)$$

By using the transformation  $x = 2\sqrt{\varepsilon}r$  we obtained the well-known standard integral of the Laguerre polynomials

$$\frac{N_{nl}^2}{(2\sqrt{\varepsilon})^{X_0/\sqrt{\varepsilon}+1}} \int_0^\infty x^{X_0/\sqrt{\varepsilon}} e^{-x} \left[ L_n^{X_0/\sqrt{\varepsilon}} (x) \right]^2 dx = 1 \quad (33)$$

The solution of the standard integral [65] is given as

$$\int_0^\infty x^{X_0/\sqrt{\varepsilon}} e^{-x} \left[ L_n^{X_0/\sqrt{\varepsilon}} (x) \right]^2 dx = \frac{\Gamma(n + X_0/\sqrt{\varepsilon} + 1)}{\Gamma(n + 1)} \quad (34)$$

Comparing equations (33) and (34) we obtained the normalization factor such that the total wave function of the mesons can be written in closed form as

$$\psi_{nl}(r) = \sqrt{\frac{(2\sqrt{\varepsilon})^{X_0/\sqrt{\varepsilon}+1} \Gamma(n + 1)}{\Gamma(n + X_0/\sqrt{\varepsilon} + 1)}} \times r^{X_0/2\sqrt{\varepsilon}} e^{-\sqrt{\varepsilon}r} L_n^{X_0/\sqrt{\varepsilon}} (2\sqrt{\varepsilon}r) \quad (35)$$

### 3. Thermal Properties of the SE with HGIQYP

To obtain the TPs of the heavy mesons, we first calculate the partition function. Equation 26 is written in the form

$$E_{nl} = Q_1 - \frac{\hbar^2}{8\mu} \left[ \frac{Q_2}{(n + \theta)} \right]^2 \quad (36)$$

where

$$\theta = \frac{1}{2} + \quad (37)$$

$$\sqrt{\left(\frac{1}{2} + l\right)^2 + \frac{2\mu}{\hbar^2 \phi^3} \left(-\frac{\vartheta R_0}{6} - R_1 \vartheta^2\right) + \frac{2\mu R_1 \vartheta^3}{\hbar^2 \phi^4} - \frac{2\mu R_1}{\hbar^2}}$$

$$Q_1 = \frac{R_0}{2} - R_1 + 2\vartheta R_1 - 2R_1 \vartheta^2 - \frac{\vartheta R_0}{2\phi} - \frac{3R_1 \vartheta^2}{\phi} + \frac{2R_1 \vartheta^3}{\phi^2} \quad (38)$$

$$Q_2 = \frac{6\mu}{\hbar^2 \phi^2} \left(-\frac{\vartheta R_0}{6} - R_1 \vartheta^2\right) + \frac{2\mu}{\hbar^2} \left(\frac{R_0}{\vartheta} + 2R_1 - 2R_1 \vartheta\right) + \frac{16\mu R_1 \vartheta^3}{3\hbar^2 \phi^3} \quad (39)$$

#### 3.1. Partition function Z(β)

The partition function (PF) takes the form [54]:

$$Z(\beta) = \sum_{n=0}^{\lambda} e^{-\beta E_{nl}} \quad (40)$$

where,  $\beta = \frac{1}{K_0 T}$ ,  $K_0$  is the Boltzmann constant,  $T$  is the absolute temperature, and  $\lambda$  is the maximum quantum number. Swapping equation (36) into equation (40) we obtain

$$Z(\beta) = \sum_{n=0}^{\lambda} e^{-\beta \left( Q_1 - \frac{\hbar^2}{8\mu} \left[ \frac{Q_2}{(n+\theta)} \right]^2 \right)} \quad (41)$$

In the classical limit, the summation is replaced by an integral,

$$Z(\beta) = \int_{\theta}^{\lambda+\theta} e^{M_1 \beta + \frac{N_1 \beta}{\rho^2}} d\rho \quad (42)$$

where

$$n + \theta = \rho \quad M_1 = -Q_1 \quad N_1 = \frac{\hbar^2 Q_2^2}{8\mu} \quad (43)$$

From equation (42), the PF is obtain as,

$$Z(\beta) = e^{M_1 \beta} \left( \frac{\rho e^{\frac{N_1 \beta}{\rho^2}} - N_1 \beta \sqrt{\pi} \operatorname{erfi} \left( \frac{\sqrt{N_1 \beta}}{\rho} \right)}{\sqrt{N_1 \beta}} \right), \quad \theta \leq \rho \leq \lambda + \theta \quad (44)$$

Other TPs can be obtained as follows:

#### 3.2. Mean energy U(β)

$$U(\beta) = -\frac{\partial}{\partial \beta} \ln Z(\beta), \quad (45)$$

#### 3.3. Free energy F(β)

$$F(\beta) = -K_0 T \ln Z(\beta) \quad (46)$$

#### 3.4. Entropy S(β)

$$S(\beta) = K_0 \ln Z(\beta) - K_0 \beta \frac{\partial}{\partial \beta} \ln Z(\beta) \quad (47)$$

#### 3.5. Specific heat capacity C(β)

$$C(\beta) = \frac{\partial U}{\partial T} = -K_0 \beta^2 \frac{\partial U}{\partial \beta} \quad (48)$$




Figure 1: Wave function of charmonium against the radius at different s state




Figure 4: Probability density function of bottomonium against the radius at different s state




Figure 2: Probability density function of charmonium against the radius at different s state




Figure 5: Partition function against  $\beta$  for different values of  $\lambda$




Figure 3: Wave function of bottomonium against the radius at different s state

#### 4. Results and discussion

The prediction of the MS of the HMs is carried out using the relation [66, 67]





$$M = 2m + E_{nl} \tag{49}$$

where  $m$  is quark-antiquark mass and  $E_{nl}$  is energy eigenvalues. Plugging equation (26) into equation (49) gives:

$$M = 2m + \frac{R_0}{2} - R_1 + 2\vartheta R_1 - 2R_1\vartheta^2 - \frac{\vartheta R_0}{2\phi} - \frac{3R_1\vartheta^2}{\phi} + \frac{2R_1\vartheta^3}{\phi^2} \tag{50}$$

$$- \frac{\hbar^2}{8\mu} \left[ \frac{\frac{6\mu}{\hbar^2\phi^2} \left( -\frac{\vartheta R_0}{6} - R_1\vartheta^2 \right) + \frac{2\mu}{\hbar^2} \left( \frac{R_0}{\vartheta} + 2R_1 - 2R_1\vartheta \right) + \frac{16\mu R_1\vartheta^3}{3\hbar^2\phi^3}}{n + \frac{1}{2} + \sqrt{\left( \frac{1}{2} + l \right)^2 + \frac{2\mu}{\hbar^2\phi^3} \left( -\frac{\vartheta R_0}{6} - R_1\vartheta^2 \right) + \frac{2\mu R_1\vartheta^3}{\hbar^2\phi^4} - \frac{2\mu R_1}{\hbar^2}} \right]^2$$

The numerical values of charmonium ( $c\bar{c}$ ) and bottomonium ( $b\bar{b}$ ), are  $m_b = 4.823 \text{ GeV}$  and  $m_c = 1.209 \text{ GeV}$ , and the corresponding reduced mass are  $\mu_b = 2.4115 \text{ GeV}$  and  $\mu_c = 0.6045 \text{ GeV}$ , respectively [68]. The potential parameters were calculated by fitting with experimental data (ED) [69].


 Figure 6: Mean energy function against  $\beta$  for different values of  $\lambda$ 

 Figure 8: Entropy against  $\beta$  for different values of  $\lambda$ 

 Figure 7: Specific heat against  $\beta$  for different values of  $\lambda$ 

 Figure 9: Free energy against  $\beta$  for different values of  $\lambda$ 

We observed that the results obtained from the prediction of MS of  $(c\bar{c})$  and  $(b\bar{b})$  mesons for different quantum states are in agreement with ED and are seen to be improved when compared with other theoretical predictions with different analytical methods from literature as shown in Tables 1 and 2.

The absolute deviation in the predicted results and ED is calculated using the following relation

$$\sigma = \frac{100}{N} \sum \left| \frac{T_P - T_{EXP.}}{T_{EXP.}} \right|, \quad (51)$$

where  $T_{EXP.}$  is the ED,  $T_P$  is the predicted results and  $N$  is the number of ED and, also the percentage error (PE) is computed with the given relation [70]

$$error = \frac{\sum ARD}{\sum T_{EXP.}} \times 100\%. \quad (52)$$

The absolute deviation was calculated using equation (51) and the values for  $(c\bar{c})$  and  $(b\bar{b})$  are 1.601 GeV and 0.457 GeV, respectively. Also, the PE of the predicted results to the ED, was calculated using equation (52) and the results show that for  $(c\bar{c})$  we have 1.60 % while that of the  $(b\bar{b})$  is 0.46 %.

In Figures 1–4 we plotted the radial wave functions and probability densities for  $(c\bar{c})$  and  $(b\bar{b})$  mesons. Generally, the probability densities decay with the radial distance but the decay happens faster at a short distance for the low-lying quantum states compare to the high-lying states. Also, the peaks of the  $(c\bar{c})$  mesons are higher than the corresponding  $(b\bar{b})$  mesons and they tend towards shorter distances. The observed high peaks may be ascribed to their heavy masses.

The variation of the PF with respect to temperature ( $\beta$ ) for different values of maximum quantum number( $\lambda$ ) is as shown

Table 1: Mass spectra of charmonium (in GeV) ( $m_c = 1.209 \text{ GeV}$ ,  $\mu = 0.6045 \text{ GeV}$ ,  $\theta = 0.06$ ,  $R_1 = -2.31507 \text{ GeV}$ ,  $R_0 = -0.15571 \text{ GeV}$ ,  $\phi = 0.67$ )

State	Our work	[1]	[64]	[44]	Experiment [69]	Absolute Relative deviation (ARD)
1S	3.096	3.096	3.096	3.095922	3.096	0.000
2S	3.686	3.686	3.686	3.685893	3.686	0.000
1P	3.327	3.214	3.433	-	3.525	0.198
2P	3.774	3.773	3.910	3.756506	3.773	0.001
3S	4.040	4.275	3.984	4.322881	4.04	0.000
4S	4.364	4.865	4.150	4.989406	4.263	0.101
1D	3.761	3.412	3.767	-	3.770	0.009
2D	4.058	-	-	-	4.159	0.101

Table 2: Mass spectra of bottomonium (in GeV) ( $m_b = 4.823 \text{ GeV}$ ,  $\mu = 2.4115 \text{ GeV}$ ,  $R_1 = -0.2542 \text{ GeV}$ ,  $R_0 = -0.37717 \text{ GeV}$ ,  $\phi = 0.67$ ,  $\theta = 0.06$ ,  $\hbar = 1$ )

State	Our work	[1]	[64]	[44]	Experiment [69]	Absolute Relative deviation (ARD)
1S	9.460	9.46	9.460	9.515194	9.460	0.000
2S	10.023	10.023	10.023	10.01801	10.023	0.000
1P	9.841	9.492	9.840	-	9.899	0.058
2P	10.160	10.038	10.16	10.09446	10.260	0.100
3S	10.345	10.585	10.28	10.44142	10.355	0.010
4S	10.522	11.148	10.42	10.85777	10.580	0.058
1D	10.142	9.551	10.14	-	10.164	0.022

in Figure 5. In Figure 5, the PF decreases as  $\lambda$  is increased. The mean energy of the mesons is depicted in Figure 6, we found that as  $\beta$  increases, the mean energy decreases monotonically. The variation of heat capacity of the mesons is presented in Figure 7. We observed an increase in the heat capacity as  $\beta$  increases for different values of  $\lambda$ , and later begin to decrease and converges at a point. The plots of the behavior of the entropy as a function of  $\beta$  is presented in Figure 8. The entropy increases to a peak value before decreasing and converges. The free energy of the mesons is plotted against  $\beta$  in Figure 9. It was noticed that the free energy decreases at different values of  $\lambda$ .

## 5. Conclusion

In this study, the HGIQYP is model to interact in quark-antiquark system and the solutions were obtained using the NU method. The energy spectrum and normalized wave function were obtained. The energy spectrum was used to predict the MS of the HMs. Also, the PF was calculated from the energy spectrum, thereafter other TPs were obtained. The results obtained showed an improvement when compared with the work of other researchers and excellently agreed with ED with a PE of 1.60 % and 0.46 % for ( $c\bar{c}$ ) and ( $b\bar{b}$ ), respectively. The PE shows that the HGIQYP is fitted in the prediction of the MS of the HMs since the PE is less.

## Acknowledgements

The Authors appreciate the referee's for the careful reading and the suggestions that improved the manuscript.

## References

- [1] H. Ciftci & H.F. Kisoglu, "Nonrelativistic-Arbitrary l-states of quarkonium through Asymptotic Iteration method", *Advances in High Energy Physics* **45** (2018) 49705.
- [2] E. Omugbe, O.E. Osafire, I. B. Okon, E.S. Eyube, E.P. Inyang, U.S. Okorie, A. Jahanshir, & C.A. Onate, "Non-relativistic bound state solutions with -deformed Kratzer-type potential using the super-symmetric WKB method: application to theoretic-information measures", *European Physical Journal D* **76** (2022) 11.
- [3] C.O.Edet, E.B.Ettah, S.A.Aljunid, R.Endut,N.Ali,A.N.Ikot, & M.Asjad, "Global Quantum information-Theoretic measures in the presence of Magnetic and Aharanov-Bohm(AB)fields", *Symmetry* **14** (2022) 976. doi.org/10.3390/sym14050976.2022.105749
- [4] E.Omugbe, O.E. Osafire, I.B. Okon, E.P. Inyang, E.S. William, & A. Jahanshir, "Any L-state energy of the spinlessSalpeter equation under the Cornell potential by the WKB Approximation method: An Application to mass spectra of mesons", *Few-Body Systems* **63** (2022) 7.
- [5] M. Allosh,Y. Mustafa, N.K. Ahmed & A.S. Mustafa, "Ground and Excited state mass spectra and properties of heavy-light mesons", *Few-Body System* **62** (2021) 26.
- [6] E. S. William, E. P. Inyang, & E. A. Thompson, "Arbitrary - solutions of the Schrödinger equation interacting with Hulthén-Hellmann potential model", *Revista Mexicana Fisica* **66** (2020) 730. https://doi.org/10.31349/RevMexFis.66.730.
- [7] E. Omugbe, "Non-relativistic eigensolutions of molecular and heavy quarkonia interacting potentials via the Nikiforov-Uvarov method", *Canadian Journal of Physics* **98** (2020)1125.
- [8] E.S. William, E.P. Inyang, I.O. Akpan, J.A. Obu, A.N. Nwachukwu, & E.P. Inyang, "Ro-vibrational energies and expectation values of selected diatomic molecules via Varshni plus modified Kratzer potential model", *Indian Journal of Physics* (2022). https://doi.org/10.1007/s12648-0222-02308-0
- [9] E.P. Inyang & E.O. Obisung, "The study of electronic states of NI and ScI molecules with screened Kratzer Potential", *East European Journal of Physics* **3** (2022) 32. https://doi.org/10.26565/2312-4334-2022-3-04.
- [10] M.C.Zhang, G.H.Sun & S.H.Dong, "Exactly complete solutions of the Schrodinger equation with a spherically harmonic oscillatory ring-shaped potential", *Physics Letters A* **374** (2010) 708.
- [11] E. S. William, E. P. Inyang, J. E. Ntibi, J.A. Obu & E. P. Inyang, "Solutions of the Non-relativistic Equation Interacting with the Varshni-

- Hellmann potential model with some selected Diatomic molecules”, *Jordan Journal of Physics* **15** (2022) 193.
- [12] M.G. Miranga, G.H. Sun & S.H. Dong, “The Solution of the second Pöschl-Teller like potential by Nikiforov-Uvarov method”, *International Journal of Modern Physics E* **19** (2010) 129.
- [13] E.P. Inyang, E.S. William, E. Omugbe, E.P. Inyang, E.A. Ibanga, F. Ayedun, I.O. Akpan & J.E. Ntibi, “Application of Eckart-Hellmann potential to study selected diatomic molecules using Nikiforov-Uvarov-Functional analysis method”, *Revista Mexicana de Física* **68** (2022) 14.
- [14] A.N. Ikot, U.S. Okorie, P.O. Amadi, C.O. Edet, G.J. Rampho, & R. Sever, “The Nikiforov-Uvarov-Functional Analysis (NUFA) Method: A new approach for solving exponential-Type potentials”, *Few-Body System* **62** (2021) 9. <https://doi.org/10.1007/s00601-021-021-01593-5>
- [15] E. P. Inyang, E.S. William, J.E. Ntibi, J.A. Obu, P.C. Iwuji, & E.P. Inyang, “Approximate solutions of the Schrödinger equation with Hulthén plus screened Kratzer potential using the Nikiforov-Uvarov-Functional analysis method: An Application to diatomic molecules”, *Canadian Journal of Physics* (2022). <https://doi.org/10.1139/cjp-2022-0030>
- [16] I.B. Okon, C.A. Onate, E. Omugbe, U.S. Okorie, C.O. Edet, A.D. Antia, J.P. Araujo, C.N. Isonguyo, M.C. Onyeaju, E.S. William, R. Horchani, & A. N. Ikot, “Aharonov-Bohm(AB) flux and thermomagnetic properties of Hellmann plus screened Kratzer potential as applied to diatomic molecules using Nikiforov-Uvarov-Functional analysis method”, *Molecular Physics* (2022) e2046295.
- [17] E. P. Inyang, E. P. Inyang, I. O. Akpan, J. E. Ntibi, & E. S. William, “Analytical solutions of the Schrödinger equation with class of Yukawa potential for a quarkonium system via series expansion method”, *European Journal of Applied Physics* **2** (2020) 36. <http://dx.doi.org/10.24018/ejphysics.2020.2.6.26>
- [18] E.P. Inyang, P.C. Iwuji, J.E. Ntibi, E.S. William & E.A. Ibanga, “Solutions of the Schrödinger equation with Hulthén-screened Kratzer potential : Application to diatomic molecules”, *East European Journal of Physics* **1** (2022) 11. <https://doi.org/10.26565/2312-4334-2022-2-02>
- [19] M. Abu-Shady, & H.M. Fath-Allah, “The effect of extended Cornell potential on heavy and heavy-light meson masses using series method”, *Journal of the Egyptian mathematical society* **23** (2019) 156.
- [20] M. Abu-Shady, T. A. Abdel-Karim, & E. M. Khokha, “Exact solution of the N-dimensional Radial Schrödinger Equation via Laplace Transformation method with the Generalized Cornell potential”, *Journal of Quantum Physics* **45** (2018) 587.
- [21] C.O. Edet, S. Mahmoud, E.P. Inyang, N. Ali, S. A. Aljunid, R. Endut, A.N. Ikot & M. Asjad, “Non-Relativistic Treatment of the 2D Electron System Interacting via Varshni-Shukla Potential Using the Asymptotic Iteration Method”, *Mathematics* **10** (2022) 2824. <https://doi.org/10.3390/math10152824>.
- [22] E. Omugbe, O.E. Osafire, & M.C. Onyeajh, “Mass spectrum of mesons via WKB Approximation method”, *Advances in High Energy Physics* **10** (2020) 1143.
- [23] E. Omugbe, O.E. Osafire, E.P. Inyang & A. Jahanshir, “Bound state solutions of the hyper-radial Klein-Gordon equation under the Deng-Fan potential by WKB and SWKB methods”, *Physica Scripta* **96** (2021) 125408
- [24] E. Omugbe, “Non-relativistic energy spectrum of the Deng-Fan Oscillator via the WKB Approximation method”, *Asian Journal of Physics and Chemistry* **26** (2020) 36.
- [25] C.A. Onate, I.B. Okon, M.C. Onyeaju & A.D. Antia, “Approximate solutions of the Schrödinger Equation for a Momentum-Dependent potential”, *Journal of Nigeria Society Physical Science* **4** (2022) 250. DOI:10.46481/jnsp.2021.209
- [26] C.A. Onate, G.O. Egharbera, & D.T. Bankole, “Eigensolution to Morse potential for Scandium and Nitrogen moniodides”, *Journal of Nigeria Society Physical Science* **3** (2021) 286. DOI: 10.46481/jnsp/2021.407
- [27] W.C. Qiang & S.H. Dong, “Proper quantization rule”, *Europhysics Letters* **89** (2010) 10003.
- [28] C.A. Onate, & M.C. Onyeaju, “Entropic system in the relativistic Klein-Gordon Particle”, *Journal of Nigeria Society Physical Science* **3** (2021) 171. DOI:10.46481/jnsp.2021.209
- [29] S.H. Dong & M. Cruz-Irisson, “Energy spectrum for a modified Rosen-Morse potential solved by proper quantization rule and its thermodynamic properties”, *Journal of Mathematical Chemistry* **50** (2012) 892.
- [30] S. H. Dong, M. Lozada-Casson, J. Yu, F.J. Angeles & A.L. Rivera, “Hidden symmetries and thermodynamic properties for a harmonic oscillator plus an inverse square potential”, *International Journal of Quantum Chemistry* **107** (2007) 371.
- [31] S. H. Dong, “Group theory approach to the Dirac equation with a Coulomb plus Scalar potential in D+1 dimensions”, *Journal of Mathematical Physics* **44** (2003) 4479.
- [32] M. Abu-shady, C.O. Edet & A.N. Ikot, “Non-relativistic Quark model under external magnetic and Aharonov-Bohm (AB) fields in the presence of Temperature-Dependent confined Cornell potential”, *Canadian Journal of Physics* (2021) 10.11139/cjp-2020-0101
- [33] R. Kumar & F. Chand, “Asymptotic study to the N-dimensional Radial Schrödinger Equation for the quark-antiquark system”, *Communication in Theoretical Physics* **59** (2013) 467.
- [34] J.P. Prasanth, K. Sebastian & V. M. Bannur, “Revisiting Cornell potential model of the Quark-Gluon plasma”, *Physica A* **558** (2020) 124921.
- [35] H. Mutuk, “Mass Spectra and Decay constants of Heavy-light Mesons: A case study of QCD sum Rules and Quark model”, *Advances in High Energy Physics* **20** (2018) 653.
- [36] H. Mutuk, “Cornell Potential: A Neural Network Approach”, *Advances in High Energy Physics* **22** (2019) 65.
- [37] A. Vega & J. Flores, “Heavy quarkonium properties from Cornell potential using variational method and supersymmetric quantum mechanics”, *Pramana-Journal of Physics* **87** (2016) 111.
- [38] R. Kumar, R.M. Singh, S.B. Bhabhar, R. Rani & F. Chand, “Analytical solutions to the Schrödinger equation for generalized Cornell potential and its application to diatomic molecules and heavy mesons”, *Modern Physics Letter A* **37** (2022) 2250010.
- [39] H. Hassanabadi, M. Ghafourian & S. Rahmani, “Study of the Heavy-Light mesons properties via the Variational method for Cornell interaction”, *Few-Body System* **15** (2016) 1040. DOI: 10.1007/s00601-015-1040-6
- [40] M. Abu-Shady, & S.Y. Ezz-Alarab, “Trigonometric Rosen-Morse Potential as a Quark-Antiquark Interaction Potential for Meson Properties in the Non-relativistic Quark Model Using EAIM”, *Few-Body Systems* **60** (2019) 66. <https://doi.org/10.1007/s00601-019-1531-y>
- [41] M. Abu-shady, H.M. Mansour & A. I. Ahmador, “Dissociation of Quarkonium in Hot and Dense media in an Anisotropic plasma in the Non-relativistic Quark model”, *Advances in High Energy Physics* **22** (2019) 615.
- [42] E.P. Inyang, E.P. Inyang, J. E. Ntibi, E. E. Ibekwe, & E. S. William, “Approximate solutions of D-dimensional Klein-Gordon equation with Yukawa potential via Nikiforov-Uvarov method”, *Indian Journal of Physics* **95** (2021) 2739. <https://doi.org/10.1007/s12648-020-01933-x>
- [43] E.P. Inyang, E.P. Inyang, E. S. William & E.E. Ibekwe, “Study on the applicability of Varshni potential to predict the mass-spectra of the Quark-Antiquark systems in a non-relativistic framework”, *Jordan Journal of Physics* **14** (2021) 345.
- [44] E.E. Ibekwe, U.S. Okorie, J.B. Emah, E.P. Inyang & S.A. Ekong, “Mass spectrum of heavy quarkonium for screened Kratzer potential (SKP) using series expansion method”, *European Physical Journal Plus* **87** (2021) 11. <https://doi.org/10.1140/epjp/s13360-021-01090-y>
- [45] K. R. Purohit, P. Jakhad & A.K. Rai, “Quarkonium spectroscopy of the linear plus modified Yukawa potential”, *Physica Scripta* **97** (2022) 044002.
- [46] S. Patel, P.C. Vinodkumar & S. Bhatnagar, “Decay rates of charmonia within a quark-antiquark confining potential”, *Chinese Physics C* **40** (2016) 053102
- [47] V. Mateu, P.G. Ortega, D.R. Entem & F. Fernandez, “Calibrating the naive Cornell model with nrqcd”, *The European Physical Journal C* **79** (2019) 323.
- [48] F. Brau & C. Sernay, “The three-dimensional Fourier grid Hamiltonian method”, *Journal of computational physics* **139** (1998) 136.
- [49] A. Bhagyesh, “Charmonium properties using the Discrete variable representation (DVR) method”, *Advances in High Energy Physics* (2021) 7. DOI:10.1155/2021/9991152
- [50] S. Jacobs, M.G. Olsson & C. Suchyta, “Comparing the Schrödinger and Spinless Salpeter equations for heavy-quark bound states”, *Physical Review D* **33** (1986) 3348.
- [51] B. Grinstein, “A modern introduction to quarkonium theory”, *International Journal of Modern Physics* **15** (2000) 461.
- [52] W. Lucha, F. Schoberl & D. Gromes. “Bound states of quarks”, *Physics Reports* **200** (1991) 240.

- [53] E. P. Inyang, A.N. Ikot, I.O. Akpan, J.E. Ntibi, E. Omugbe, & E. S. William, "Analytic study of thermal properties and masses of heavy mesons with quarkonium potential", *Result in Physics* (2022) 105754. <https://doi.org/10.1016/j.rinp.2022.105754>
- [54] M. Abu-Shady , T. A. Abdel-Karim & Y. Ezz-Alarab, "Masses and thermodynamic properties of heavy mesons in the non-relativistic quark model using the Nikiforov-Uvarov method", *Journal of Egyptian Mathematical Society* **23** (2019)155.
- [55] E.P.Inyang,E.P.Inyang, I.O.Akpan, J.E.Ntibi, & E.S.William, "Masses and thermodynamic properties of a Quarkonium system", *Canadian Journal Physics* **99** (2021) 990.<https://doi.org/10.1139/cjp-2020-0578>.
- [56] X. Cao, S. Qiu, H. Liu & D. Li, "Thermal properties of light mesons from holography" **8** (2021) 005.
- [57] E.P. Inyang, E.S. William, J.O. Obu, B. I. Ita , E.P. Inyang & I.O. Akpan, "Energy spectra and expectation values of selected diatomic molecules through the solutions of Klein-Gordon equation with Eckart-Hellmann potential model", *Molecular Physics* **119** (2021) e1956615 <https://doi.org/10.1080/00268976.2021.1956615>
- [58] I.B Okon,E. Omugbe, A.D. Antia, C. A. Onate, L.E. Akpabio, & O.E. Osafire, "Spin and pseudospin solutions to Dirac equation and its thermodynamic properties using hyperbolic Hulthén plus hyperbolic exponential inversely quadratic potential", *Scientific Reports* **11** (2021)21.
- [59] P.O. Okoi, C.O. Edet, T.O. Magu & E.P. Inyang, "Eigensolution and Expectation values of the Hulthén and Generalized Inverse Quadratic Yukawa potential", *Jordan Journal of Physics* **15** (2022) 148.
- [60] L. Hulthén, "Über die eigenlösungen der Schrödinger-Gleichung des deuterons", *Ark. Mat. Astron. Fys. A* **28** (1942) 5 .
- [61] O. Oluwadare & K. Oyewumi "Non-Relativistic Treatment of a Generalized Inverse Quadratic Yukawa Potential", *Chinese Physics Letters* **34** (2017) 110301.
- [62] S.K. Nikiforov, & V.B. Uvarov. *Special functions of Mathematical Physics* (Birkhauser, Basel, 1988).
- [63] E. P. Inyang, E. S. William & J.A.Obu, "Eigensolutions of the N-dimensional Schrödinger equation interacting with Varshni-Hulthén potential model", *Revista Mexicana de Fisica* **67** (2021) 193. <https://doi.org/10.31349/RevMexFis.67.193>
- [64] M. Abu-Shady, "N-dimensional Schrödinger equation at finite temperature using the Nikiforov-Uvarov method", *Journal of Egyptian Mathematical Society* **21** (2016) 4.
- [65] I.S. Gradshteyn & I.M. Ryzhik, *Tables of integrals, series and product*, 7th ed. Academic Press, Boston (2007).
- [66] I.O. Akpan, E.P. Inyang, E.P. Inyang & E.S. William, "Approximate solutions of the Schrödinger equation with Hulthén-Hellmann Potentials for a Quarkonium system", *Revista Mexica De Fisica* **67** (2021) 490.
- [67] E. P. Inyang, J. E. Ntibi, E.A. Ibang, F. Ayedun, E. P. Inyang, E.E. Ibekwe, E. S. William & I.O. Akpan, "Thermodynamic properties and mass spectra of a quarkonium system with Ultra Generalized Exponential- Hyperbolic potential", *Communication in Physical Science* **7** (2021)114.
- [68] R. Olive, D. E. Groom & T.G. Trippe, "Particle Data Group", *Chinese Journal of Physics C* **38** (2014) 60.
- [69] M. Tanabashi , C. D. Carone,T.G.Trippe, & C.G. Wohl, "Particle Data Group", *Physical Review D* **98** (2018) 546.
- [70] E.P. Inyang, P.C. Iwuji, J.E. Ntibi, E. Omugbe, E.A. Ibang & E.S. William, "Quark-antiquark study with inversely quadratic Yukawa potential using Nikiforov-Uvarov-Functional analysis method", *East European Journal of Physics* **2** (2022) 51. <https://doi.org/10.26565/2312-4334-2022-2-05>